

Projet de Recherche INTERREG-V océan Indien 2014-2020
A1/OT1/OS-01a - Action II-2 TN

ReNovRisk-Cyclones et Précipitations



L6: Rapport ou article scientifique sur l'analyse préliminaire des résultats obtenus lors de la campagne expérimentale de 2019

Olivier BOUSQUET

Laboratoire de l'Atmosphère et des Cyclones
(UMR 8105 CNRS/Météo-France/Université de La Réunion)



Livrable 6

Rapport ou article scientifique sur l'analyse préliminaire des résultats obtenus lors de la campagne expérimentale de 2019

Contexte

La campagne expérimentale du projet RNR-CP visait à échantillonner les propriétés des systèmes dépressionnaires tropicaux dans la région du Canal du Mozambique lors de la saison cyclonique 2018-2019 et à former des étudiants et personnels techniques de la région dans le domaine de l'observation météorologique. Les opérations menées dans le cadre du projet ont été combinées avec d'autres, réalisées au cours de la même période, dans le cadre du projet RNR-C3.

Les opérations ont essentiellement consisté en une campagne de radiosondages atmosphériques qui s'est déroulée du 20 janvier au 8 avril 2019. Ces activités ont été mises à profit pour former des dizaines d'étudiants et chercheurs malgaches, mahorais et mozambicains tout au long des 3 mois de campagne. Les trois sites de radiosondages étaient situés à Maputo (site de l'INAM), Mayotte (site de la DIROI) et Tamatave (site de l'ASECNA - aéroport).

Une autre opération, consistant à déployer en parallèle des instruments d'observation de l'OPAR sur Petite Terre (Mayotte) a dû être annulée au vu du contexte social très préoccupant qui régnait à Mayotte en 2018. Considérant les risques importants de dégradation et de retard lié à l'envoi d'instruments sur place, les instruments ont été opérés depuis La Réunion.

Résultats

Les travaux de recherche basés sur l'exploitation de ces observations ont donné lieu à la rédaction de deux articles scientifique dédiés, tous deux publiés dans la revue *Atmosphere* (Impact Factor 3.11).

Le premier (Bousquet et al. 2021) est un article de synthèse proposant un bilan de toutes les activités réalisées au cours du programme dans le domaine de l'observation :

Bousquet O, Barruol G, Cordier E, Barthe C, Bielli S, Calmer R, Rindraharisaona E, Roberts G, Tulet P, Amelie V, Fleischer-Dogley F, Mavume A, Zucule J, Zakariasy L, Razafindradina B, Bonnardot F, Singh M, Lees E, Durand J, Mekies D, Claeys M, Pianezze J, Thompson C, Tsai C-L, Husson R, Mouche A, Ciccione S, Cattiaux J, Chauvin F, Marquestaut N. Impact of Tropical Cyclones on Inhabited Areas of the SWIO Basin at Present and Future Horizons. Part 1: Overview and Observing Component of the Research Project RENOVRIK-CYCLONE. *Atmosphere*. 2021; 12(5):544. <https://doi.org/10.3390/atmos12050544>

Le second (Durand et al. 2021) est un article centré sur l'exploitation des mesures réalisées depuis l'île de La Réunion :

Durand J, Lees E, Bousquet O, Delanoë J, Bonnardot F. Cloud Radar Observations of Diurnal and Seasonal Cloudiness over Reunion Island. *Atmosphere*. 2021; 12(7):868. <https://doi.org/10.3390/atmos12070868>

Ces deux articles sont joints ci-après



Article

Impact of Tropical Cyclones on Inhabited Areas of the SWIO Basin at Present and Future Horizons. Part 1: Overview and Observing Component of the Research Project RENOVRIISK-CYCLONE

Olivier Bousquet, Guilhem Barruol, Emmanuel Cordier, Christelle Barthe, Soline Bielli, Radiance Calmer, Elisa Rindraharisaona, Gregory Roberts, Pierre Tulet, Vincent Amelie et al.

Special Issue

Tropical Cyclones in the Indian Ocean







Edited by

Prof. Dr. Olivier Bousquet



Article

Impact of Tropical Cyclones on Inhabited Areas of the SWIO Basin at Present and Future Horizons. Part 1: Overview and Observing Component of the Research Project RENOVRISK-CYCLONE

Olivier Bousquet ^{1,2,*}, Guilhem Barruol ³, Emmanuel Cordier ⁴, Christelle Barthe ^{1,5} , Soline Bielli ¹, Radiance Calmer ^{1,6}, Elisa Rindraharisaona ^{3,7} , Gregory Roberts ^{8,9}, Pierre Tulet ^{1,5}, Vincent Amelie ¹⁰, Frauke Fleischer-Dogley ¹¹, Alberto Mavume ¹² , Jonas Zucule ¹³, Lova Zakariasy ¹⁴, Bruno Razafindradina ¹⁴, François Bonnardot ¹⁵, Manvendra Singh ¹⁶, Edouard Lees ¹, Jonathan Durand ¹, Dominique Mekies ¹, Marine Claeys ^{1,9}, Joris Pianezze ¹, Callum Thompson ¹, Chia-Lun Tsai ^{1,17} , Romain Husson ¹⁸, Alexis Mouche ¹⁹ , Stephane Ciccione ²⁰, Julien Cattiaux ⁹, Fabrice Chauvin ⁹  and Nicolas Marquestaut ^{1,4}



Citation: Bousquet, O.; Barruol, G.; Cordier, E.; Barthe, C.; Bielli, S.; Calmer, R.; Rindraharisaona, E.; Roberts, G.; Tulet, P.; Amelie, V.; et al. Impact of Tropical Cyclones on Inhabited Areas of the SWIO Basin at Present and Future Horizons. Part 1: Overview and Observing Component of the Research Project RENOVRISK-CYCLONE. *Atmosphere* **2021**, *12*, 544. <https://doi.org/10.3390/atmos12050544>

Academic Editor: Corene Matyas

Received: 31 March 2021

Accepted: 20 April 2021

Published: 23 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

- ¹ Laboratoire de l'Atmosphère et des Cyclones (UMR8105 LACy), Université de La Réunion, CNRS, Météo-France, 97400 Saint-Denis, France; christelle.barthe@aero.obs-mip.fr (C.B.); soline.bielli-bousquet@univ-reunion.fr (S.B.); radiance.calmer@colorado.edu (R.C.); pierre.tulet@aero.obs-mip.fr (P.T.); edouard.lees@meteo.fr (E.L.); jonathan.durand@live.com (J.D.); dominique.mekies@meteo.fr (D.M.); marine.claeys@meteo.fr (M.C.); joris.pianezze@univ-reunion.fr (J.P.); callum-thompson@ucsb.edu (C.T.); chialun@knu.ac.kr (C.-L.T.); nicolas.marquestaut@univ-reunion.fr (N.M.)
 - ² Institute for Coastal Marine Research (CMR), Nelson Mandela University, Port-Elizabeth 6031, South Africa
 - ³ Institut de Physique du Globe de Paris, Université de Paris, CNRS, 75238 Paris, France; barruol@ipgg.fr (G.B.); elisa.rindraharisaona@univ-reunion.fr (E.R.)
 - ⁴ Observatoire de Sciences, Univers de La Réunion (UMS 3365 OSU-R), 97400 Saint-Denis, France; emmanuel.cordier@univ-reunion.fr
 - ⁵ Laboratoire d'Aérodynamique, Université de Toulouse, UT3, CNRS, IRD, 31400 Toulouse, France
 - ⁶ Cooperative Institute for Research in Environmental Sciences, National Snow and Ice Data Center (NSIDC), University of Colorado Boulder, Boulder, CO 80304, USA
 - ⁷ Laboratoire GéoSciences Réunion (LGSR), Université de La Réunion, 97400 Saint-Denis, France
 - ⁸ Scripps Institution of Oceanography, University of California, San Diego, CA 92093, USA; greg.roberts@meteo.fr
 - ⁹ Centre National de Recherche Météorologique (UMR3589 CNRM), Université de Toulouse, CNRS, Météo-France, 31057 Toulouse, France; julien.cattiaux@meteo.fr (J.C.); fabrice.chauvin@meteo.fr (F.C.)
 - ¹⁰ Seychelles Meteorological Authority, International Airport, Mahe 670311, Seychelles; v.amelie@meteo.gov.sc
 - ¹¹ Seychelles Islands Foundation, Mont-Fleuri, POB 853, Victoria, Mahe 670311, Seychelles; ceo@sif.sc
 - ¹² Department of Physics-Faculty of Sciences, Eduardo Mondlane University, Maputo CP 257, Mozambique; amavume@yahoo.co.uk
 - ¹³ Instituto Nacional de Meteorologia (INAM), Maputo CP 256, Mozambique; jonas_z@inam.gov.mz
 - ¹⁴ Institut Supérieur de Technologie d'Antsiranana, Antsiranana BP 509, Madagascar; lova.zakariasy@ist-antsiranana.mg (L.Z.); hbrazafindradina@ist-antsiranana.mg (B.R.)
 - ¹⁵ Direction Interrégionale de Météo-France pour l'Océan Indien, 97400 Saint-Denis, France; francois.bonnardot@meteo.fr
 - ¹⁶ Mauritius Oceanography Institute, Morcellement de Chazal, Albion 95410, Mauritius; msingh@moi.intnet.mu
 - ¹⁷ Department of Astronomy and Atmospheric Sciences, Center for Atmospheric REmote Sensing (CARE), Kyungpook National University, Daegu 41566, Korea
 - ¹⁸ Collecte Localisation Satellites (CLS), 29280 Brest, France; rhusson@cls.fr
 - ¹⁹ Laboratoire d'Océanographie Physique et Spatiale, Ifremer, CNRS, IRD, UBO, 29280 Plouzané, France; Alexis.Mouche@ifremer.fr
 - ²⁰ Kelonia, Observatoire des Tortues Marines de La Réunion, 97436 Saint-Leu, France; stephane.ciccione@museesreunion.re
- * Correspondence: olivier.bousquet@meteo.fr

Abstract: The international research program “ReNovRisk-CYCLONE” (RNR-CYC, 2017–2021) directly involves 20 partners from 5 countries of the south-west Indian-Ocean. It aims at improving the observation and modelling of tropical cyclones in the south-west Indian Ocean, as well as to foster regional cooperation and improve public policies adapted to present and future tropical cyclones risk in this cyclonic basin. This paper describes the structure and main objectives of this ambitious

research project, with emphasis on its observing components, which allowed integrating numbers of innovative atmospheric and oceanic observations (sea-turtle borne and seismic data, unmanned airborne system, ocean gliders), as well as combining standard and original methods (radiosoundings and global navigation satellite system (GNSS) atmospheric soundings, seismic and in-situ swell sampling, drone and satellite imaging) to support research on tropical cyclones from the local to the basin-scale.

Keywords: tropical cyclone; south-west Indian Ocean; gliders; unmanned airborne system; bi-logging; sea turtles; global satellite navigation system; ReNovRisk; numerical modelling; climate modelling; austral and cyclonic swells; seismic data

1. Introduction

Due to their highly destructive potential, tropical cyclones (TCs) have long been considered a major risk for populations, territorial economies, and biodiversity. In this regard, predicting their outcome and impacts at present and future times is one of the major concerns of both the Intergovernmental Panel on Climate Change (IPCC) and the World Meteorological Organization (WMO).

As highlighted in the latest reports of WMO's International Workshop on Tropical Cyclones (IWTC), research carried out over the last four years has considerably improved our understanding of TC intensification processes [1,2], as well as TC tracking and intensity forecasting [3,4]. The operational implementation of coupled ocean-atmosphere (OA) numerical weather prediction (NWP) systems by many national weather services has, in particular, played a key role in reducing forecasting errors at all space and time scales [5–8]. Despite these important advances, additional efforts are still needed to accurately predict and characterize the potential impacts of tropical cyclones on a given territory, especially during landfall. Such efforts include, for instance, the collection of novel atmospheric and oceanic observations, to better constrain (and verify) the performance of coupled NWP systems [9,10], as well as the implementation of wave models and specific microphysical parameterizations to improve roughness, swell, wind speed, and momentum flux representation in TC forecasting systems [11–13].

Accurate modelling of OA interactions is particularly crucial in areas such as the tropical south-west Indian Ocean (SWIO) basin (30–90° E, 0–40° S), where the atmospheric variability is associated with a particularly strong oceanic response (and vice versa). The SWIO (Figure 1), which contributes to approximately 10–12% of the worldwide cyclonic activity [14–16], is indeed widely considered as the cyclonic basin with the highest prevalence of OA interactions [17] due to the unique structure of the thermocline in the Seychelles-Chagos Thermocline Ridge area (55–70° E, 5–15° S) [18,19]. Like most TC basins, the SWIO includes many fragile countries, whose economic development and infrastructure, as well as food, medicine and water supply chains, are regularly impacted by tropical cyclones.




Figure 1. Map of the south-west Indian Ocean (SWIO) tropical cyclone basin (30–90° E, 0–40° S). The principal locations discussed in the paper are indicated by dark blue (France), light blue (Mozambique), green (Mauritius), red (Madagascar), and pink (Seychelles) circles.

In very recent years, countries bordering the Mozambique Channel (MC) have indeed been struck by a series of extremely intense and devastating events, whose economic impact will be felt for many years to come. Heavy rains associated with TC DINEO (2017) have caused 700,000 refugees and tens of millions of USD of damage in Mozambique, while the overall cost of TCs ENAWO (2017) and AVA (2018), which affected nearly one million people in Madagascar, was estimated to be more than USD 600 million (about 7% of Madagascar's average annual gross domestic product). These heavy tolls are, however, out of all proportion to those of TCs IDAÏ (considered by the United Nations as the worst natural disaster ever in the MC) and KENNETH (the most intense TC ever reported in the MC [20]), which both made landfall in Mozambique in 2019 [21]. According to the latest economic reports, these two storms have affected a total of nearly 1.7 million people and caused damage and losses estimated at ~USD three billion—plus a further recovery cost estimated at USD 3.4 billion—by the World Bank and Mozambican officials [22].

Given the colossal impact of TCs on the local populations, infrastructure, and economic development of many countries in the SWIO basin, the European Union (EU), together with the Regional Council of Réunion Island and the French State, have designed the transdisciplinary research program "Réunion NOVative research on cyclonic RISks" (ReNovRisk), to improve the resilience of SWIO countries to TC hazards and mitigate associated economic vulnerability, damages, and risks (e.g., winds, rainfall, landslides, submersion) in inhabited areas. To achieve these objectives, ReNovRisk has been divided into four interlinked research projects, referred to as ReNovRisk-Cyclone, -Erosion, -Impacts and -Transfer, whose overall objectives are described in [23]. The present paper focuses on the Cyclone component of this program, which involves a large international consortium of research institutes, universities, and weather services originating from France (e.g., universities of Réunion Island and Toulouse, Centre National de Recherche Scientifique (CNRS), Météo-France, Institut de Physique de Globe de Paris (IPGP), Institut National de l'Information Géographique et Forestière (IGN), Institut Français de Recherche pour l'Exploitation de la MER (IFREMER)), Mozambique (Eduardo Mondlane and Pemba Unilurio universities, Mozambique Weather Service (INAM)), Madagascar (Institut Supérieur de Technologie de Diego Suarez, Université d'Antananarivo), the Seychelles (Seychelles Meteorological Authority, Seychelles Islands Foundation), Mauritius (Mauritius Oceanography Institute), as well as international institutions such as the European Space Agency (ESA) and WMO, among others.

Through its observing, modelling, climate and outreach components, ReNovRisk-Cyclone (hereafter referred to as RNR-CYC) aims to improve the observation and modelling of TCs, as well as to provide inputs for other components of the global ReNovRisk program focusing on hydrological (RNR-Erosion and RNR-Transfer) and economical (RNR-Impact)

consequences of tropical cyclones [23]. Another key objective of RNR-CYC is to foster regional cooperation and improve public policies adapted to present and future TC risks faced by territories bordering the SWIO. The latter is all the more essential because predicted changes in the coupled OA system due to global warming are likely to generate significant modifications of the cyclonic activity in the coming decades. Consequently, regions that are currently spared or moderately affected by TCs, and that often lack experience-based adaptation strategies, may soon have to face potentially increasing TC-related hazards [24,25]. Such changes include, for instance, the widening of the tropical belt resulting from ocean warming [26,27], which has already been shown to induce a poleward migration of TC's lifetime maximum intensity (LMI) in both hemispheres [28–31], or significant modifications in TC frequency and/or length of the TC season [32].

This paper aims to describing the structure and main objectives of the project RNR-CYC, as well as to present an overview of the main applications and results of its observation component—modelling aspects are presented in more details in the companion paper [33]. This article is organized as follows: Section 2 provides an overview of the four components of RNR-CYC (observation, mesoscale modelling, climate modelling, cooperation, and outreach). Section 3 presents the major achievements of RNR-CYC regarding oceanic and atmospheric observations, while Section 4 concludes and discusses new research topics to be investigated beyond the end of this program.

2. Structure and Objectives of RNR-CYC

The project RNR-CYC focuses on the meteorological and oceanographic impacts of TCs in the SWIO (Figure 1) at both present and future horizons. It aims, in particular, to better apprehend the impacts of these extreme storms on the main inhabited islands of this oceanic basin by providing innovative modelling and observing products that will also feed the cascade risk analysis tools deployed in the other research components of the global ReNovRisk program [23]. In order to achieve these objectives, RNR-CYC has been divided into four components (Figure 2): (i) an observation component, to improve both long-term and temporary observations of TCs and their atmospheric and oceanic environments; (ii) a mesoscale modelling component, to improve modelling and short-term forecasting of TCs; (iii) a climate component, to evaluate the consequences of climate change on the variability and structure of TCs at both local and basin scales; and (iv) an outreach component, aimed at improving capacity building in the three aforementioned research areas through strengthening cooperation between SWIO countries. An overview of the structure and objectives of these four components is described hereafter.



Figure 2. Structure and main objectives of ReNovRisk-CYCLONE.

2.1. Observing Component

The observing component of RNR-CYC is aimed at providing additional observations of TCs and their environment by improving regional and local observing capabilities in the SWIO. It has been built around three complementary approaches:

(i) A “conventional” approach, based on the reinforcement of regional ground-based meteorological observation facilities and, in particular, of the water vapor Global Navigation Satellite System (GNSS) observation network operated by the International GNSS Service (IGS). Starting in November 2017, 10 new public observation sites (composed of ground-based GNSS receivers and collocated surface weather stations) have been deployed in Madagascar, Eparses Islands, and the Seychelles in the frame of RNR-CYC’s sub-program “Indian Ocean GNSS Applications for Meteorology” (IOGA⁴MET [34]) to increase the number of tropospheric GNSS measurements (e.g., zenithal delay, integrated water vapor amounts) and positioning data throughout the western part of the basin (see Section 3.2.2.). The first analysis of GNSS-derived observations collected during RNR-CYC have already demonstrated the benefit of these new permanent stations to investigate the water vapor cycle at diurnal to inter-annual time scales [34,35], but also to provide new and continuous observations to investigate the dynamics of the Earth’s crust in this particularly active part of the world [34,36].

(ii) An “experimental” approach, based on the temporary collection of atmospheric and oceanic observations at various locations in the basin. For this purpose, several atmospheric and oceanographic field campaigns have been organized throughout the 3.5-year duration of RNR-CYC, with the goal to provide novel datasets to evaluate numerical model developments and simulations performed in the frame of the project [33]. The main achievements include:

- A three-year satellite acquisition campaign (2017–2020), set up in collaboration with ESA and IFREMER, to collect high-resolution (1 km) observations of surface winds and sea roughness from spaceborne synthetic aperture radars (SARs) deployed onboard the satellites Sentinel 1A/1B of the European Earth Observation Program Copernicus (<https://www.copernicus.eu/fr>, see Section 3.3., accessed on 20 March 2021);
- A regional field campaign, organized from late January to early April 2019, to investigate atmospheric and oceanic environmental conditions prevailing in the vicinity of TCs during the 2018–2019 TC season. During this 2.5 month period, a regional radiosounding network, allowing for the collection of nearly 500 soundings, was deployed in Mayotte (France), Toamasina (Madagascar) and Maputo (Mozambique) to both sample the atmospheric environment of TC and train students and academics in experimental meteorology (see Section 2.4.);
- The deployment of two ocean gliders from Réunion Island to sample the vertical properties of the upper ocean layers in the Mascarene Archipelago (see Section 3.1.4.);
- The deployment of an unmanned airborne system (UAS), equipped with aerosol, turbulence, sea state, and meteorological sensors to measure OA fluxes and aerosol concentrations off the shore of Réunion Island (see Section 3.2.1.);
- The organization of several local observation campaigns to sample sea swell properties during austral winters and summers using acoustic Doppler current profilers (ADCP) and wave gauges deployed near the shore of Réunion Island (see Section 3.1.1.).

(iii) An “exploratory” approach, based on the deployment and evaluation of innovative methods to collect oceanographic observations. A particularly original approach, based on biologging technology, has been evaluated for two years to collect data from sea turtles (ST) equipped with dedicated ARGOS environmental tags in the frame of RNR-CYC’s subprogram “Sea Turtle for Ocean Research and Monitoring” (STORM, see Section 3.1.3.). Another original approach, based on the previous work of [37–39], was also further investigated to quantify extreme swell phenomena from microseismic noise measurements recorded by ground seismometers (see Section 3.1.2.). The preliminary assessment of terrestrial seismic observations collected in Réunion Island against oceanographic records and offshore wave model data have demonstrated that land-based seismic stations could be particularly useful to observe both austral [40] and cyclonic swell [41] (this Special Issue).

2.2. Modelling Component

Protecting life and property requires a precise estimate of the environmental changes associated with the passage of TCs in the vicinity of inhabited areas. The challenge in the face of the cyclonic threat is to simultaneously predict the track and intensity of the storms, but also the consequences resulting from their landfall, or transit near inhabited areas. Hence, damages caused to a given territory, which are essentially related to rainfall intensity, wind strength and sea state (e.g., swell), could significantly differ depending on whether it is affected by a tropical storm (TS), a monsoon depression, or a more or less intense TC. To this end, many operational meteorological services and research centers concerned with TC hazards have made considerable efforts to develop deterministic and ensemble coupled NWP systems providing high spatial resolution forecasts in all TC basins [42–46]). Another fundamental element to improve TC impact predictions, which is partly addressed in the hydrological component (RNR-Erosion) of the global ReNovRisk program, is to also take into account the role of land surface and hydrological processes even after the dissipation of the storm. The recent case of TC IDAÏ, whose catastrophic damage in Beira was mainly caused by flooding occurring days after the storm had passed over the city, is a perfect example of the urgent need to also improve hydrological forecasting.

Improving TC forecasting first and foremost implies a proper representation of the interactions between the storm and the ocean, and vice versa [47]. During the propagation of a TC over an oceanic area, mixing caused by surface winds usually induces a significant drop in surface temperature [48,49] that strongly reduces surface enthalpy and heat fluxes [50,51]. These air–sea fluxes can also be significantly impacted by waves, which redistribute momentum in the near-surface layer and modify the enthalpy fluxes through the emission of sea spray [52–57]. In this regard, new parameterizations reproducing the impact of marine aerosols on turbulent heat exchanges have been proposed and validated in recent years [58,59], but are yet to be implemented in atmospheric models to thoroughly evaluate their impact on TC behavior.

Radiative cooling [60,61], evaporation [62,63] and latent heat release [64] have long been recognized to play a key role in the development and intensification of tropical cyclones; therefore, particular attention must also be paid to microphysical schemes implemented in NWP systems. These schemes must, in particular, allow for an efficient representation of the radiative cooling at the top of the storms (which is a constraining criterion of TC intensity) and of the vertical distribution of latent heat (which represents the main source of energy of TCs). They should thus also be able to realistically take into account the role of atmospheric aerosols, that (indirectly) affect the radiation balance by impacting on the radiative and precipitating properties of the clouds. Improving TC forecasting therefore also implies the development of coupled aerosol–microphysical–radiation schemes to be integrated in fully coupled ocean–wave–atmosphere (OWA) models.

An important objective of RNR-CYC was to develop high-resolution OWA and OA modelling systems capable of representing, as exhaustively as possible, the multitude of physical interactions that control the variations of intensity of TCs, as well as their impacts (wind, rain, swell) at the scale of SWIO territories. The main modelling developments made in the frame of this project can be found in [11,65], (this Special Issue, [66]) and in the companion paper [33].

2.3. Climate Component

Evaluating the impact of climate change on the frequency and intensity of tropical cyclones is considered as one of the top five issues of concern by the IPCC. Currently, regional and global climate models make it possible to identify the preferred areas of cyclogenesis and occurrence of tropical low-pressure systems at the basin scale, but cannot yet be effectively relied upon to investigate potential changes in their structure and intensification mechanisms. Although it is now widely accepted that the global increase in sea surface temperatures in tropical areas will be a favorable element for TC development, it is not clear how other ingredients involved in their formation and intensification will evolve in

the future. In this regard, another important objective of RNR-CYC was to evaluate the global evolution of cyclonic activity in the SWIO, but also to investigate potential structural and intensity changes of TCs resulting from the ongoing modification of their oceanic and atmospheric environments.

The modelling strategy was based on two complementary approaches, allowing to both estimate the evolution of cyclonic activity at the basin scale (i.e., changes in trajectory, intensity, and frequency of TCs at different time scales), and to assess potential structural changes and impacts of TCs at the local scale. This strategy relies on the exploitation of unprecedented high-resolution global climate simulations [32], as well as of mesoscale coupled simulations to estimate the impact of climate change on the intensity, behavior, and consequences of TCs at the scale of a given territory. Examples of results obtained from such high-resolution model runs are discussed in [33,67] (both in this Special Issue).

2.4. Regional Cooperation

Another important objective of RNR-CYC is to provide enhanced tools and knowledge to SWIO countries facing cyclonic hazards. Responding to this strong societal issue requires a better structuring of the regional scientific community, as well as significant reinforcement of the cooperation between countries bordering the SWIO basin. While regional collaboration already exists through WMO's regional structures (e.g., Tropical Cyclone Programme (<https://public.wmo.int/en/our-mandate/focus-areas/natural-hazards-and-disaster-risk-reduction/tropical-cyclones>, accessed on 20 March 2021) and Réunion Island's Regional Specialized Meteorological Center (RSMC La Réunion, http://www.meteo.fr/temps/domtom/La_Reunion/webcmrs9.0/anglais/index.html, accessed on 20 March 2021)) and the Indian Ocean Committee (<https://www.commissionoceanindien.org/>, accessed on 20 March 2021), interactions remain essentially focused on operational and technical applications, and only modestly promote research development in this area.

One of the strengths of RNR-CYC consists of the implementation of a partnership involving many regional research institutes, universities, and meteorological services that agreed to pool their resources and expertise to strengthen the resilience to TC hazards, and develop public policies better adapted to the risks faced by SWIO territories. RNR-CYC is thus a fundamentally collaborative project that is not only based on the large sharing of data and experiences, but also on training programs and capacity-building initiatives in the fields of observation and forecasting. These actions include, for instance, the organization of forecasting training sessions at RSMC La Réunion, of training courses in climatology and climate change, as well as numerous co-supervised MSc internships based on the analysis of experimental measurements collected in the project.

For instance, the project's field phase, which involved nearly one hundred participants from late January to mid-April 2019, was an opportunity for many students and researchers to initiate themselves to the technique of atmospheric radiosounding (RS). During this 2.5-month campaign, ~500 radiosoundings were performed from three experimental sites specifically deployed for this occasion in Maputo (INAM's headquarters, Mozambique), Mayotte (Météo-France weather center, France), and Toamasina (Toamasina international airport, Madagascar). On this occasion, nearly 60 students and academics from Antananarivo (Madagascar), Eduardo Mondlane (Mozambique) and Réunion universities came to Toamasina and Maputo to conduct RS measurements, while many senior forecasters of INAM and Météo-France Mayotte had, for the first time, the opportunity to operate and familiarize themselves with an RS station (Figure 3).




Figure 3. The Toamasina (Madagascar) radiosounding campaign. (a) Training of students from Antananarivo University at Toamasina airport. (b) Statistical analysis of the 140 RS measurements performed in Toamasina: mean altitude of the tropopause, ascending speed, and maximum RS altitude. Picture: Olivier Bousquet

The involvement of the French consular services in the project also made it possible to communicate widely to the general public, the scientific community, and the media in Mozambique, Madagascar and the Seychelles on the issues of adaptation to climate change and natural hazards. As will be seen in Section 4, these regional collaborations will continue for many years to come through several new research projects initiated by RNR-CYC.

3. Results

3.1. Oceanic Observations

3.1.1. In Situ Swell Observations

One of the main objectives of RNR-CYC is to assess the impact of tropical cyclones on land, including possible submersion resulting from cyclonic swell surge along coastlines. In many tropical islands, the latter are bordered by coral reefs that can serve as a defense against flooding. These reef systems, particularly fringing reefs, protect the coastline by acting as low-pass filters that can reduce the energy of wave flows reaching the coast by up to 98% in the gravitational part of the wave spectrum [68]. The physical processes underlying coral reef coastal protection consist of a complex combination of incident waves, tides, and wind-induced surges [69–71].

Ocean wave energy is concentrated in the gravitational frequency band of the wave spectrum, generally between 0.04 Hz and 0.25 Hz. These gravitational waves (GWs) are the main drivers of the hydrodynamics of reef systems, as well as of beaching, runup, and submersion. During and after breaking at the edge of reef systems, GWs are dissipated while low frequency waves (infragravity waves ($0.004 < IG \text{ (Hz)} < 0.04$) and very low-frequency waves ($0.001 < VLF \text{ (Hz)} < 0.004$) propagate to the shore. Previous analyses of these propagation and transformation processes across various coral reefs have shown that wave dynamics could vary considerably depending on the characteristics and location of reef systems [71,72].

In order to both quantify the physical processes linked to severe sea states and assess the protective role of reef systems, a cross-shore transect, composed of bottom-fixed wave gauges and ADCPs, was deployed in Réunion Island from February to April in 2019 and 2020. Instruments were installed at the fringing reef of “Trou d’Eau”, located along the west coast of the island (Figure 4a). At the near-shore site, reef-base (RS) and reef-flat (RF) stations were deployed through the fringing reef along a cross-shore transect. The RS station was installed at the base of the reef slope at an average depth of 12 m, while three RF stations (RF1, RF2 and RF3 in Figure 4a) were aligned on the flat reef inside the lagoon at a depth of 1 m (Figure 4b,c). RS and RF stations were all equipped with synchronized ocean sensor system instrument (OSSSI) wave gauges, enabling continuous recording pressure at a sampling frequency of 10 Hz. The RS station also featured an upward-looking Nortek AQuadopp (AQP) profiler configured to measure current profiles every 20 min, with a 2 Hz hourly burst mode to record wave parameters. In 2020, an ADCP (RDI Sentinel V100)

was also deployed at an ocean offshore (OC) site at a depth of 45 m. This instrument was configured to record incident wave parameters from hourly bursts of 2100 samples at 2 Hz, and current profiles from the bottom to the surface.




Figure 4. Overview of the experimental setup deployed in “Trou d’Eau” (Réunion Island) in February–April 2019 and 2020. (a) Coastal study area, (b,c) details of the cross-shore transect instrumented during the experiments. The stations are labeled OC for ocean offshore, RS for reef-slope, and RF for reef-flat. In (c), labels OSSI, AQP and RDI ADCP correspond to wave gauges, Nortek AQuadopp current Profiler, and RDI ADCP current profiler, respectively. Instrument deployment dates are indicated in (c).

Observations collected at the offshore stations OC and RS were relied upon to describe and quantify the main properties (height, period, and direction) of the waves impinging on the reef in the GW frequency band (wave periods ranging from 4 to 25 s), while data gathered at RF stations were used to investigate wave transformation (across the fringing reef) and propagation (from outside the lagoon to the shore). Observations collected at OC and RS stations were processed with the RDI software “Velocity” and the PUV method [73], respectively. All OSSI pressure data were corrected from atmospheric mean sea level pressure data recorded at the surface weather station of “La Rivière Des Galets” (located 20 km northward of the experimental site, RDG on Figure 4a) and non-hydrostatic pressure following the linear wave theory [74].

The mean surface elevation was computed from a 20 min moving average of the signal, while the wave spectrum and average wave parameters were obtained through the application of a fast Fourier-transform (FFT). The latter was applied over 4096 data points for incident gravity wave band or swells ($0.04 < SW < 0.25$ Hz), and 32,768 data values for infragravity ($0.004 < IG < 0.04$ Hz) and very low frequency ($0.001 < VLF < 0.004$ Hz) bands.

The recorded signal was not perfectly periodic; therefore, a Hamming window, leading to zero value at the edges, was also applied to mitigate artifacts resulting from leakage issues. To avoid resulting loss of information at the edges, a 30-min (resp. 3 h) time average was then performed for the incident (resp. IG and VLF) bands. Time series of power spectral density deduced from data collected in 2019 along the cross-shore transect are shown in Figure 5, together with the associated mean power spectra at each station.




Figure 5. Spectral wave characteristics observed along the instrumental transect presented Figure 4b. The plots are organized in the shoreward direction from the top to the bottom (reef slope RS1 station on the top, and reef flat station RF1 closest to the shore on the bottom). The left panels show the temporal variation of the wave spectrum recorded at each station from February to April 2019. The black solid line indicates the frequency cut at 0.04 Hz between the gravity frequency band (GW) and the infragravity (IG) frequency band. The right panels show the mean power spectral density for each station averaged over the whole period, with the frequencies band GW, IG and VLF (for very low frequency) indicated.

Measurements collected at the reef slope station RS clearly show that the wave energy spectrum is concentrated in the gravity band, with an averaged peak period of 13.5 s over the whole period. After breaking at the reef crest, most of this energy is dissipated (by breaking and/or by friction on the reef bottom), while the remaining part (in IG and VLF bands) is transferred inside the reef system. For the two strong wave events observed in February (likely associated with TC GELENA to the north of Réunion Island) and April 2019 (austral swell event), the total wave energy reduction between the reef slope station RS and the adjacent reef flat station RF3 reaches 98%, a value in good agreement with previous meta-analyses [68]. Within the reef, the transfer of the remaining energy between the three stations was nevertheless slightly different for these two events. In February (TC event), the wave energy reduction reached 64% from RF3 to RF2 and 2% from RF2 to RF1, but only 47% from RF3 to RF2 and 15% from RF2 to RF1 in April (austral swell event). The main difference between these two events is related to the period of the incident waves (11.8 s for the TC-related event and 16 s for the austral swell event); this result suggests that the

generation and propagation of IG waves inside the reef might be less important for short waves, induced by the wind, than for longer-period swells. This shoreward propagation with less energy reduction is also noticeable in the average spectra, which show a small translation of the peak frequency toward higher frequencies in the IG band, between the RF2 and RF1 station, and a reduction in the peak amplitude.

These preliminary analyses show that Réunion Island's fringing reef plays a strong role in protecting the shore against incident waves. Further investigations will be conducted in order to thoroughly investigate the physical processes and the role of both reef topography and roughness on the dynamics of IG waves. This includes, for instance, the impact of shoreward-propagating IG waves on onshore suspended-sediment transport [75], and the relationship between long wave propagation (and transformation) across the reef and sea water level above the reef—the remaining long waves propagating through the reef have been shown to increase the back-reef set-up and beach runup, which could lead to increased coastal erosion during extreme events such as tropical cyclones or strong storms [76,77].

3.1.2. Ground-Based Swell Observations

The global monitoring of swell activity induced by tropical storms (TS) and TCs is of major interest to quantify the risk associated with extreme swells, but also to validate numerical models used to predict ocean activity. Direct swell observations such as those presented in the previous section are, however, strongly limited by the low number of oceanographic sensors available in this area, as well as by their deployment (and servicing) costs and their vulnerability during tropical cyclones. These limitations motivated the use of indirect observations as alternative and complementary observables to quantify the swell parameters. In this regard, the analysis of wave-induced seismic noise is known to be an interesting substitute for monitoring ocean activity and has been proven to be particularly relevant for assessing the impact of waves on coastal environments [38]. The possibility to derive swell measurements from the seismic noise generated by ocean swell and transmitted to the solid earth as seismic waves recorded by terrestrial seismological instruments [78], is discussed hereafter from data collected in RNR-CYC.

Microseisms recorded by seismic stations worldwide are known to be generated by ocean gravity waves [79]. Seismic energy spectra at terrestrial seismic stations showed two clear peaks in separate frequency bands, known as primary and secondary microseisms (hereafter named PM and SM, respectively), originating from different physical processes involving local or distant sources of ocean wave activity [80]. Primary microseisms (PM) are mostly visible at coastal and island stations and are assumed to be generated by direct interactions of swell-induced pressure variation with the coastal seafloor [81–83]. PM noise has the same periods as the ocean swell (i.e., between 8 and 20 s), providing a powerful way to characterize the local impact of swell approaching the shore. On the other hand, SM noise is generally generated in the deep oceans and at larger distances from coastal areas [37,84–86]. It dominates seismic noise at both continental and oceanic stations and exhibits a large peak at half the period of ocean waves (i.e., between 3 and 10 s), generated by the interference of swells of similar periods travelling in opposite directions [79] and generating seismic surface waves travelling horizontally within the solid crust.

In the Indian Ocean, most remote sources of seismic noise are located in the southernmost part of the Austral Ocean basin and are associated with storm systems moving around Antarctica [84,87,88]. Some noise sources may also develop at tropical latitudes in association with tropical cyclones [41] (this Special Issue). Recent seismic deployments on the ocean floor enabled the making of in situ observations of SM underneath TCs in the neighborhood of Réunion Island [37] and confirmed the possibility to track TC and TS from the ocean bottom. Although SM are generally created by distant storms, they can also be generated by the coastal reflection of waves if incident and reflected waves propagate in opposite directions [89,90]. In this latter case, the incoming swell may interfere with its reflected swell, resulting in the generation of standing waves close to coastal areas that

oscillate at twice the frequency of the propagating wave [91]. Some studies suggest that local and distant sources of noise in the SM frequency peak may coexist [39,92,93].

Previous seismic analyses conducted in the Pacific and Indian Oceans have already demonstrated that several swell parameters can be derived from the seismic data. This includes: (i) the swell peak period T_p , derived from the dominant frequency of the PM and SM using the power spectral density analysis of the seismic records; (ii) the local or distant wave significant height H_s (in the case of the PM or SM band, respectively), obtained by measuring the microseism amplitudes through hourly Root Mean Square (RMS); and (iii) the wave peak direction D_p for the case of the PM band, or the source direction in the case of the SM band, which can both be deduced from the polarization analysis of the three seismic data components, to determine the dominant direction and strength of the recorded microseismic noise.

To illustrate this innovative approach, we analyse hereafter seismic observations collected during the tropical storm (TS) ELIAKIM that developed in March 2018. This storm has clear signatures, despite not reaching tropical cyclone intensity, and developed at a large distance from Réunion Island, demonstrating the potential of the method in quantifying remote systems. As mapped on Figure 6a, TS ELIAKIM started as a depression located north-east of Madagascar on 13 March 2018. It then intensified to a tropical storm on 14 March and to a strong tropical storm on 15 and 16 March while approaching the eastern coast of Madagascar, where it made landfall on 17 March. TS ELIAKIM continued its southward motion and returned over the ocean on 18 March. After a final burst on 19 March, the storm collapsed on 20 March, while entering its extratropical transition phase.

Our analysis was performed from data recorded by seismic stations of the permanent seismic network of the Piton de la Fournaise Volcano Observatory (OVPE, seismic code PF, red triangles on Figure 6b) and from a temporary seismic network deployed in the frame of the RNR program (code ZF, blue triangles on Figure 6b). Seismic data were compared with swell parameters issued from the WaveWatch3 (WW3) model [94,95] at nodes surrounding the island from the global wave model hindcast [96], indicated by stars on Figure 6b.

The seismological energy content of the vertical ground displacement of station MAT (Figure 6b) is shown in the spectrogram of Figure 7a. It displays the temporal variation of the Power Spectral Density (PSD) during the period 10 to 25 March 2018 over the frequency band of the ocean activity (0.05–0.5 Hz, i.e., 20–2 s periods). Superimposed to the spectrogram is the distance curve of the storm center to Réunion Island (dashed line), together with the storm intensity curve (black line indicating the mean wind speed and the colored dots the storm classification, as in Figure 6a). The seismic energy at this station shows good correlation with the storm intensity, despite its large distance, varying from 500 to 1500 km with two pulses of energy with the maximum at a frequency range of 0.1–0.3 Hz, i.e., in the SM band, culminating on 16 and 19 March during maximum storm intensity. Below 0.1 Hz, i.e., in the PM band, the PSD still displays clear energy increasing during the two storm peaks.




Figure 6. (a) Track of TS ELIAKIM as derived from RSMC La Réunion best-track data. The colored circles indicate the position and intensity of the storm every 6 h. (b) Locations of the seismic stations (blue triangle, temporary ZF network, and red triangle, permanent PF stations) and WaveWatch3 nodes [96] surrounding the island, where wave height H_s (colored stars) is extracted. Vectors indicate the average azimuth on March 19, 2018, obtained from the polarization analysis in the SM frequency band (green) and computed from the storm track and the RER seismic station (pink).

The hourly RMS amplitudes of the seismic noise recorded at the various seismic stations on the island are shown in Figure 7b,c, for both the SM and the PM bands, respectively. Note that the vertical axes are different, and that the SM amplitude is almost one order of magnitude larger than the PM. The 21 seismic stations analyzed over Réunion island display similar variation patterns. This indicates that this noise is not a purely local source; the storm acting as a distant SM source, or that the source—if local—is larger than the size of the island, which is the case for the swell generating the PM. The observed station-to-station variability in RMS amplitudes suggests the influence of local site effects such as the installation, the coupling of the seismic sensor with the ground, the nature of the bedrock, and the attenuation around the station.

The RMS peaks at seismic station MAT (Figure 7b) are observed on 16 and 19 March. This period corresponds to the maximum intensity of TS ELIAKIM and to the local variation of the swell height, as visible on H_s data issued from the WW3 model at various points around Réunion island (Figure 8). The maximum H_s observed on 19 March (corresponding to the nodes at the longitude of 55° E) interacted with the local bathymetry slightly later, which explains the small delay with the PM. A small peak was also observed on 21 March in both SM and PM data with dominating energy at period ~10s, likely originating from a distant source and not related to any local swell activity increase.




Figure 7. Temporal variation of the microseismic noise recorded during TS ELIAKIM. (a) Spectrogram at seismic station MAT (see location in Figure 6b) between 10 and 25 March and up to 0.5 Hz, together with the storm intensity (continuous black line) and distance between the storm center and the seismic station (dashed black line). The colored circles indicate the intensity of the storm every 6 h, as on Figure 6a). (b) Secondary microseisms RMS amplitude variations measured at the island seismic stations (left axis, in micrometers), together with the distance between the storm and seismic station (right axis, in km). (c) PM RMS amplitude variations (left axis) at Réunion Island seismic stations.

If one accepts that the SM is generated in the vicinity of the storm center and that the PM is generated in coastal areas closer to the seismic stations, one should observe a delay between the two curves. Such a delay was previously observed for austral swells generated by distant storms [40] and proposed as a precursor for predicting coastal submersion in Réunion island. In the present case, considering the involved distances (TS ELIAKIM was located ~ 800 km of the coast of Réunion Island on 16 March, and 500 km on 19 March) and the involved velocities (3 km s^{-1} for the surface waves carrying the SM signal, and $\sim 50 \text{ km h}^{-1}$ for the long period waves at the surface of the ocean), one should expect a delay of >10 h for each peak. However, Figure 8 shows a delay of only ~ 4 h between the SM and PM. This may indicate that the SM is not generated at the storm center, but likely closer to the island. Alternatively, it may also indicate a slow wind–wave growth that may take from a few hours to few days [97]. The swell–noise amplitude correlation can be used to build a transfer function to translate the amplitude of the seismic noise in terms of swell height. Although such a relation is station-related, and therefore not universal, it was nevertheless shown to provide particularly good results for strong swell events [40,41,81].




Figure 8. Seismic and swell amplitudes recorded during TS ELIAKIM in March 2018. SM (black dots) and PM (pink dots) RMS amplitude at seismic station MAT (see location Figure 6b) versus the significant wave heights (H_s) at different nodes around La Réunion (colored lines). Note that the amplitude of the PM is multiplied by 8 to reach about the same scale as the SM. The green and magenta dots indicate the maximum SM and PM, respectively, showing a delay of 4 h between the two. The significant waves heights H_s extracted at the various model nodes around the island (Figure 6b) are plotted in continuous colored lines with the same color codes as the stars in Figure 6b.

The recordings of the three components of the ground motion also make the seismic noise a vectorial observation for measuring the direction and strength of the signal polarization. In the case of the SM band, the polarization is expected to point towards the noise source, i.e., the storm center, whereas it should indicate the very local swell propagation direction in the case of the PM. In this PM case, some angle may exist between the swell propagation direction at a large distance offshore and the swell at the coast, due to refraction of the swell and the coastal bathymetry. In the case of TS ELIAKIM, the polarization was measured on an hourly basis in the SM frequency band. Figure 6b displays the average polarization recorded on 19 March at the various land seismic stations in the 0.1–0.33 Hz frequency band (pink) that can be compared to the azimuth computed from the storm center position. This map illustrates a very homogeneous orientation of the noise polarization across the island, pointing to the SW (toward the storm center location on that date) and confirming that the SM originated in the vicinity of the storm center.

These results confirm that seismic noise may provide a useful proxy to quantify the swell parameters. In the PM band (10–20 s period), the seismic noise amplitude allows for a robust quantification of the swell height H_s through a transfer function, the spectrogram allows deciphering the swell dominant period T_p , and the polarization analysis allows retrieving the local swell direction D_p . In the SM band (3–10 s period), the amplitude revealed the storm strength and the polarization indicated the storm azimuth. Terrestrial seismic stations therefore provide alternative and complementary observations of both TC and ocean activities. In some cases, the availability of several decades of seismic archives may also provide new opportunities to derive cyclone climatologies [41].

3.1.3. Biologging Observations

Observing the vertical structure of the ocean is essential to improve knowledge of both the coupled OA system and marine ecosystems. In this regard, an increasingly common alternative to gather high-resolution hydrographic profiles in the world's oceans is to rely on animal-borne sensors (a.k.a. biologging) to collect in situ observations in remote and under-instrumented areas. Compared to conventional oceanographic in situ observation

approaches (e.g., gliders, ARGO drifters, buoys, research cruises), animal-borne electronic ARGOS tags are relatively inexpensive to operate and can be deployed in remote areas with limited human resources. This approach offers all countries the possibility to actively contribute to the collection of ocean observations; therefore, biologging is expected to grow considerably in the future. The recent decision, in August 2020, of the Global Ocean Observing System (GOOS)'s Executive Committee to create a new observing network exclusively dedicated to animal-borne ocean sensors ("ANIBOS") is definitely in line with this perspective and clearly attests of the immense potential of this approach.

The potential of biologging for sampling the thermal structure of the tropical Indian Ocean was evaluated in the frame of RNR-CYC's sub-program "Sea Turtle for Ocean Research and Monitoring" (STORM). STORM was initiated in January 2019 by Réunion's University and Réunion Island's Sea-Turtle Observatory (Kelonia), with the goal of monitoring the state of the tropical Indian Ocean down to several hundred meters below the surface in near-real-time and at high spatial (<100 m) and temporal (5') resolutions. As of March 2021, 22 animals have been equipped with Temperature–Depth (TD) ARGOS tags before being released from Réunion Island (Figure 9). Note that all animal manipulations were made by accredited ST biologists and that all STs were accidentally captured in fisherman's nets in the vicinity of Réunion Island and brought back to Kelonia's care center to be healed and rehabilitated.




Figure 9. Trajectories and names of the 18 loggerheads and Olive Ridley STs equipped with Wildlife Computers SPLASH10 tags released from Réunion's Island as of 10 March 2021. Circles next to the ST names on the right-hand side indicate active tags as of 20 March 2021. The insert on the left-hand side shows a zoomed view of ST Union trajectory between 28 August and 17 December 2020. Four additional animals, equipped with LOTEK Kiwisat tags, were also released from Réunion island in 2019 (not shown). Plots are extracted from Wildlife Computers' real-time ARGOS tracking platform.

During this experiment, two species of late juvenile sea turtles (loggerhead and Olive Ridley) were equipped with Argos TD tags. While some animals stayed in their pelagic habitats, some loggerheads also began their first homing migration to the Oman Gulf, thus enabling the collection of data both in tropical areas, from loggerheads, and subtropical areas, from Olive Ridelys (Figure 9). STs released from Réunion Island principally swam at (or slightly below) the surface (~50% of the time) and near the bottom of the ocean mixed layer (OML, ~25% of the time). They were found to dive up to 100 times a day, sometimes up to 350 m, allowing the collection of numerous hydrographic profiles within and far below the OML [9]. The analysis of data collected during the first year of this experiment has confirmed the great potential of this approach for sampling the vertical structure of

the ocean, validating ocean models and spaceborne sensors, as well as to investigate the intra-seasonal variability of the tropical Indian Ocean [9].

Sea turtles are known to generally swim in rings and frontal areas in between, and often travel by moving from one ring to another [98]. These rings and eddies are numerous in the Mozambique Channel [99,100], but less common north of Réunion Island, which may explain that loggerhead STs released from there generally tend to follow a straight trajectory (Figure 9)—another explanation (currently under investigation) is that STs rarely feed during their reproductive migration, so as to reach their breeding areas as fast as possible. Nevertheless, according to its circular trajectories, ST Union (Olive Ridley), which moved south-east of Réunion Island in late 2020, apparently travelled in oceanic gyres and eddies (Figure 9).

The surface current analysis of Mercator Ocean’s operational model NEMO-PSY4 (a.k.a Glo12, [101]), averaged from 15 October to 15 December, confirmed that ST Union indeed swam in cyclonic and anticyclonic eddies of variable sizes during this period (Figure 10a). Vertical cross-sections of the ocean temperature field across the center of the main eddy (Figure 10b,c) show strong upwellings at the edges of the vortex (while downwelling can be observed at the center), resulting in significant temperature gradients at the surface (Figure 10a). This vertical transport, which brings nutrient-rich waters from the thermocline up to the surface, makes these eddies prime feeding areas for many marine species and are particularly appreciated by sea turtles. Equipping STs with environmental tags is thus an easy and relatively affordable way to sample the properties of these important transient mesoscale features, which will be further investigated in the continuation of the STORM program (see Section 4).




Figure 10. GLO12 model analyses in the area of evolution of ST Union averaged from 15 October to 15 December 2020. (a) surface currents and sea surface temperature superimposed on ST Union track over the 2 month period (green dots), (b,c) vertical cross-sections of ocean temperature through the center of the main vortex travelled by ST Union.

During this experiment, three sea turtles were also caught in the immediate vicinity of tropical cyclones during TC seasons 2018–2019 and 2019–2020: ST Brice, which moved in the vicinity of TC KENNETH (April 2019) during its cyclogenesis (Figure 11a), and STs India and Tom that were trapped in TC HEROLD (March 2020) during its intensification phase (Figure 11b). Spaceborne oceanic observations are generally unavailable under

cloudy conditions; therefore, data collected by these animals also represent a fantastic opportunity to investigate the impact of TCs on the surface and subsurface structure of the ocean.

Sea surface temperature data collected by ST India in TC HEROLD from 14 to 20 March 2020 are shown in Figure 11c. During these six days, the animal remained trapped in the immediate vicinity of the storm center (~ 30 km from the TC eye) and only moved ~50 km in the north/north-west direction. In situ surface temperature observations collected from 14 to 17 March showed SST cooling of ~3.5°C (29 °C to 25.5 °C) in 72 h—during these three days, the animal remained quasi-stationary. As the storm progressively moved south-eastwards, ST India began to slowly move to the north-west, over the area previously affected by the tropical cyclone. Observations collected from 17 to 20 March showed that the temperature surface layer in this area quickly returned to pre-storm conditions, to regain its initial temperature of 29 °C on 19 March. As shown in [33], surface and subsurface observations collected by STs in tropical cyclones can also provide key data to evaluate coupled model forecasts in cyclonic conditions.




Figure 11. Satellite images of (a) TC KENNETH on 15 April 2019 and (b) TC HEROLD on 15 March 2020. Sea turtle symbols show the location of ST Brice during the cyclogenesis of TC KENNETH (left panel) and of STs India and Tom during the intensification phase of TC HEROLD (right panel). (c) Evolution of sea surface temperature (°C) in the vicinity of TC HEROLD, as measured by ST India between 14 and 20 March 2020 within the area (51.93–52.62° E; 13.9–14.67° S).

3.1.4. Glider Observations

On 22 January 2019, two SlocumG1 gliders operated by CNRS were deployed from Réunion Island for a period of two months. The two instruments were programmed to follow a north-east (glider GLNE) and a north-west (glider GLNW) trajectory to reach the northernmost region of the Mascarene Archipelago, where a high probability of TC formation was suggested about 400–600 km north of Réunion Island (Figure 12). The data acquisition strategy was set in order to complete a saw-tooth navigation pattern, allowing the gliders to dive with an angle of 26° between 5 m and 980 m depth (resulting in an along-track resolution of about 4 km once the profile was normalized on the vertical).




Figure 12. Trajectories of the two gliders deployed from Réunion Island between 22 January 2019 and 22 March 2019. (a) Trajectories of glider GLNE (blue) and GLNW (red), together with the trajectory and intensity (colored dots) of TC GELENA between 5 and 9 February. The positions of the gliders during the TC period are indicated by black ellipses. Right panels show cast depth range and mean current velocities along the glider tracks estimated from glider drift for (b) GLNE and (c) GLNW. Table 1: Glider instrumentation, sampling rate (Hz), vertical resolution (m) and depth range (m) of collected data.

The two gliders were programmed to sample the ocean during descending (downcast) and ascending (upcast) profiles. Observations were transmitted in real time by satellite telemetry after each upcast, when directives for modifying the sampling strategy and glider navigation (based on operational ocean model forecasts) were also received. Each glider was equipped with various physical and optical biogeochemical instruments for sampling the ocean temperature, salinity, oxygen, turbidity, and chlorophyll-a concentration at different rates according to depth, as shown in Table 1. All sensors were operational for glider GLNW, but the oxygen and optical sensors were turned off on glider GLNE.

The primary objective of this experiment was to investigate OA kinetic energy exchanges in cyclonic conditions, with emphasis on the fraction of kinetic energy transmitted to the ocean. As shown by [102], who analyzed the vertical structure of the ocean in a tropical cyclone sampled during the CIRENE field phase [103], this kinetic energy is generated on the left side of the TC track in the southern hemisphere, and later consumed by strong vertical mixing resulting from surface water cooling [104]. Due to the presence of strong currents ($>0.25 \text{ m s}^{-1}$) along the glider trajectories, an important surface drift prevented the full application of the initially planned northward navigation strategy. The navigation parameters were then modified early in the mission to mitigate the battery consumption by reducing the depth range of the glider (because the oil pump that controls the glider buoyancy is the device that consumes the most energy). The gliders profiles were thus reduced to 500 m and 300 m depths (as shown by the colored tracks on Figure 12b,c). Real-time data transmission was also turned-off at some points to reduce the time spent at the surface, when drifting was maximized. The slower-than-expected displacement speed of the gliders prevented them from intercepting the core of TC GELENA, which developed in the area between 5 and 9 February 2019 (Figure 12a). Thanks to real-time trajectory optimization, both systems were nevertheless able to approach relatively close to the storm, as shown by black ellipses displayed in Figure 12a. For instance, glider GLNE was able to move to a distance of about 290 km from the TC center and to collect ocean data in the direct vicinity of this storm.

Table 1. Glider instrumentation, sampling rate (Hz), vertical resolution (m) and depth range (m) of collected data.

Parameter	Instrument	Sampling	Resolution	Depth
GLNE				
Temperature Salinity and Depth	CTD Seabird SBE-41cp	1/8	1.5	−5 to −980
Oxygen	Aanderaa Optode 5013	-	-	-
Fluorescence Turbidity	Wetlabs flbbcd	-	-	-
GLNW				
Temperature Salinity and Depth	CTD Seabird SBE-41cp	1/8	1.5	−5 to −980
Oxygen	Aanderaa Optode 4831	1/8	1.5	−5 to −980
Fluorescence Turbidity	Wetlabs flbbcd	1/8	1.5	−5 to −980

Vertical profiles of temperature, salinity and density collected by the two gliders between 0 and 150 m and 23 January and 17 February 2019 (along the two northward segments of the tracks) are shown in Figure 13, together with corresponding mean profiles for each parameter. The mixed layer depth (MLD), indicated by a black solid line, is estimated as the depth where the temperature differs by 0.2 °C from a surface reference value of 10 dbar [105,106]. Due to the significant distance separating the two gliders from the TC core, and because the maximum kinetic energy generated in the upper ocean occurred on the left side of the TC track, no significant change in the surface temperature could be noticed during the pre-cyclonic, cyclonic, and post-cyclonic phases (Figure 13a,d,g). A relatively constant temperature of 29 °C was observed between the surface and 30 to 40 m depth, but quickly decreased to reach 22 °C at a depth of 150 m. Salinity measurements (Figure 13b,e,h) indicated a freshening of the water within the top 100 m layer, starting around 6 February (−19.22° N) for the GLNW track, and from 28 January (−20.15° N) for the GLNE track. While part of this freshening can be explained by the rainfall generated by TC GELENA between 15° S and 5° N, another possible explanation is related to the advection of fresher water originating from the southern branch of the South Equatorial Current, which flows westward with a 20 Sv transport to feed the East Madagascar Current [107]. The velocity field computed from Mercator-Ocean’s global reanalysis model PHY-001-030, which indicated the presence of strong westward currents between 28 January and 9 February (not shown), and the vertical extent of this freshwater lens, both seem to support this hypothesis. Density measurements (Figure 13c,f,i) did not seem to be impacted by this freshening and showed a well-mixed layer within the top 40 m of the ocean and a well-stratified layer underneath.

The mixed layer depth shows significant frequency variations in relation to the diurnal cycle, and also appears deeper along the GLNW track until 8 February. GLNW was always located southward of GLNE before this date; therefore, these observations suggest a sloping up of the MLD in the northward direction. As GLNW moved northward of GLNE after 8 February, the MLD measured by GLNE became deeper, thus reinforcing the hypothesis of a northward sloping-up of MLD. A significant rise of the MLD of up to 20 m (on 5 and 6 February for GLNE and GLNW, respectively) could also be observed almost immediately after the TC formation (4 February). The MLD then returned to its initial value between 9 February (for GLNW) and 12 February (for GLNE).




Figure 13. Time series of vertical temperature (top), salinity (middle), and density (bottom) profiles within the top 150 m ocean layer from 23 January 2019 to 17 February 2019 for (a–c) GLNW and (d–f) GLNE tracks. Panels (g–i) show the associated mean vertical profiles over the same period. The vertical white dashed line in (a–f) indicates the date of formation of the Tropical Cyclone GELENA (4 February). The vertical blue dashed line shows the closest position of each glider to the TC core (around 9 February). The top axis indicates the latitudinal location of the profiles.

The temperature–salinity (TS) diagrams collected along the full glider tracks (23 January to 23 March 2019) are shown in Figure 14. The water masses sampled by the two instruments, known as the Indian Central Water, were formed and subducted in the Subtropical Convergence area of the southern gyre of the Indian Ocean [107,108]. The OML in this water mass was generally between 50 and 150 m in depth, with temperature between 25 and 30 °C and salinity in the range of 34.9–35.4 psu. Observations collected from the bottom of the OML to 1000 m showed salinity values between 34.5 and 36 psu and temperatures decreasing from 25 °C to 5 °C.




Figure 14. Temperature–salinity (TS) diagram along (a) GLNW and (b) GLNE tracks from 23 January to 23 March 2019. Inserts show corresponding glider tracks.

3.2. Atmospheric Observations

3.2.1. UAS Observations

During RNC-CYC field phase, a UAS system called “BOREAL” was operated from Réunion Island to sample the lower part of the atmosphere (Figure 15). The main objective of this experiment was to provide in situ measurements of air–sea interactions in cyclonic and pre-cyclonic conditions to evaluate numerical models and microphysical parameterizations developed in the frame of the program.

The BOREAL UAS (Boréal SAS, Toulouse, France) is a fixed-wing aircraft with a 4.2 m wingspan propelled by a thermal engine (Figure 15b) that has a maximum take-off weight of 25 kg and can fly in winds up to 25 m s^{-1} . For this experiment, a 5 kg scientific payload was developed to study air–sea interactions using an optical particle counter for measuring the aerosol number and size distribution ($0.3 < \text{diameter} < 3.0 \text{ }\mu\text{m}$; MetOne), a custom-designed multi-hole probe for measurements of turbulence and three-dimensional winds, a radar altimeter for wave height and sea state measurements, a broadband shortwave pyranometer for downwelling solar radiation (Licor), an infra-red temperature sensor to measure sea-surface temperature, as well as standard meteorological measurements (atmospheric pressure, temperature and relative humidity). With this payload, the endurance and range of the BOREAL UAS was $\sim 7 \text{ h}$, 700 km, and 2000 m above sea level (asl). A live video was also streamed up to 40 km from the ground-station to provide additional safety during low-altitude segments when the UAS flew $\sim 40 \text{ m}$ asl. The BOREAL UAS is autonomous, although its flight plan could be adapted at any time to accommodate weather conditions or air traffic via a radio or a satellite link. A transponder mounted on its wing also allowed it to integrate into international airspace.

The BOREAL UAS flew over the Indian Ocean in two exclusive zones allocated by the French aviation authority to the southeast and northwest sides of Réunion Island (Figure 15a). These two areas were dedicated for scientific flights over the international waters up to 250 km from the ground station and with a ceiling at 1067 m above sea level. BOREAL UAS operations were conducted from the airfields of Cambaie (when flying towards the north of the island) and Bras Panon (when flying towards the south). In total, 12 scientific flights were carried out between February and March 2019, for a total of 53.7 h of research flights and 5012 km of observations over the ocean. Three flights of more than 200 km from the ground station were carried out, the longest of which lasted 6 h and 18 min and covered 610 km. Examples of observations collected by the BOREAL UAS during this two-month operating period are presented below.




Figure 15. (a) The 12 BOREAL flights operated from Réunion Island during the RNR-CYC campaign in February and March 2019. The yellow and orange polygons represent the authorized flight zones to the northwest and southeast of Réunion Island. The outer red circle denotes an operating radius of 250 km around Réunion Island. (b) The BOREAL UAS leaving the catapult for a scientific mission from Cambaie, Réunion Island. Sources: (a) Google Earth; (b) Greg Roberts.

Figure 16 summarizes the BOREAL UAS measurements of aerosol concentrations for particle diameters $>0.3 \mu\text{m}$ and $>1.0 \mu\text{m}$ within 200 m above the ocean surface, compared to horizontal wind speed (Figure 16a) and wave height (Figure 16b) for conditions encountered during the two-month campaign. As expected, there was an increase in particle concentrations (diameter $>0.3 \mu\text{m}$), which is often associated with primary marine aerosol (PMA) emissions, over the observed range of wind speed (2.2 to 13.5 m s^{-1}) and wave heights (2 to 3.7 m). However, the range of wind speeds and wave heights encountered in the vicinity of Réunion Island during the RNR-CYC field campaign remained relatively small, partly because the paths of the TCs were never closer than a few hundred kilometers to the ground station during the observed period.




Figure 16. Aerosol concentrations (diameter $>0.3 \mu\text{m}$ and $>1.0 \mu\text{m}$) measured during the BOREAL flights as a function of (a) the averaged horizontal wind in the marine boundary layer and (b) the measured wave height.

Although strong cyclonic conditions were not encountered in the vicinity of Réunion Island during the 2019–2020 TC season, several BOREAL UAS flights were nevertheless impacted by the presence of TC JOANINHA, which developed in the eastern part of the Mascarene Archipelago in March 2019 (Figure 17a). The signature of this storm could be seen up to several hundred kilometers away in observations of ocean waves, atmospheric turbulence structures, and enhanced PMA emissions. These emissions, which occur over a much larger domain than the cyclonic system, have been shown to modify the TC's track and intensity [11].

As TC JOANINHA moved to the east of Mauritius, the BOREAL UAS flew to the west of Réunion Island to sample the TC's perimeter (see flight track in Figure 17a and wave crests, swells and clouds generated by the cyclone recorded by the on-board camera in Figure 17b). In situ measurements were used, in particular, to assess parameterizations of air–sea interactions simulated with the OWA coupled system developed in RNR-CYC (see [33] for details about this system). Results presented hereafter are derived from a coupled OWA simulation based on the ocean model Coastal and Regional Ocean Community (CROCO, <http://www.croco-ocean.org>, accessed on 20 March 2021), the wave model Wave Watch 3 (WW3) [94,95], and the atmospheric model Meso-NH (<http://mesonh.aero.obs-mip.fr/>, accessed on 20 March 2021), which was developed specifically for comparisons with the BOREAL UAS.

Figure 18 presents a vertical profile of aerosol particles conducted during the 25 March flight from near the ocean surface ($<50 \text{ m asl}$) to above the marine boundary layer ($\sim 1000 \text{ m asl}$). Observed aerosol concentrations (diameter $>0.3 \mu\text{m}$) were found to be relatively constant throughout the marine boundary layer (up to $\sim 800 \text{ m asl}$) and decreased above the inversion. A surface layer with enhanced emissions, similar to the profiles observed at the surf zone in the frame of project Miriad [109], was also observed at the lowest part of the vertical profile ($\sim 50 \text{ m asl}$).




Figure 17. (a) Satellite image of TC JOANINHA on 25 March 2019 in the vicinity of Mauritius and Réunion Island. (b) picture from the BOREAL on-board camera showing the ocean sea-state and the TC on the horizon. Sources: (a) MSG satellite; (b) Christophe Mazel

This surface layer, which was not captured in the model simulation (Figure 18), has not been well-documented so far and has important implications for the transport of aerosols into the well-mixed boundary layer. In addition, the simulations tend to overestimate the aerosol emissions near the ocean surface (i.e., an approximate factor of two enhancement in aerosol concentrations) and show a pronounced vertical gradient throughout the marine boundary layer with an underestimate of aerosol concentrations in the free troposphere. These large differences between observed and simulated vertical profiles of aerosol concentrations suggest that mixing timescales in the marine boundary layer may perhaps be too slow in the model—resulting in an excessive gradient in the aerosol concentration—but could also be related to a possible bias in the CAMS/COPERNICUS analyses used to initialize sea salts in Meso-NH. Although a mass distribution calculation was made to adjust the consistency of aerosol distributions in CAMS and Meso-NH, a small remaining mass difference in the finest modes can also lead to large differences in the number of aerosols, especially in areas with low primary aerosol production, i.e., where aerosol number concentrations mostly arise from initial conditions.




Figure 18. BOREAL profile of aerosol numbers for particles larger than 0.3 μm obtained with the onboard Optical Particle Counter (green) and simulated with the OWA model (orange).

A few days later, a southern swell event occurred when winds shifted to the north-east direction as TC JOANINHA continued its eastward progression through the Indian Ocean. Flight operations were moved to Bras Panon and the BOREAL UAS flew along a southern curtain for more than 200 km to the east of Réunion Island on 29 March 2019 from 6:33 to 12:04 UTC (Figure 19a). The platform followed a flight plan with ascents from 100 to 1000 m asl followed by straight-level legs at 400 and 100 m asl for at least 10 km each. Mean and significant wave height observations collected during this flight by the onboard radar altimeter showed that wave height increased by about one meter (from ~2.5 to 3.5 m) between locations A and D (Figure 19b).




Figure 19. (a) WW3 simulation of significant wave height on 29 March 2019 with the BOREAL UAS flight overlaid on the image (black line). The blue crosses correspond to A, B, C and D legs of the flight at 100 m asl. (b) Significant wave height (H_{m0}) and mean wave height from the BOREAL UAS observations compared to wave height simulated by WW3.

A similar increase in wave height was also obtained from the simulations of this case study (Figure 19a). The associated modelled (Figure 20a) and observed (Figure 20b) spectral density functions provide insight on the composition of the southern swell and show two peaks for both observations and simulations. Azimuth plots from WW3 based on the wave elevation spectrum (Figure 20c) indicate that these peaks corresponded to distinct components of the southern swell; the first component was generated by the northerly winds (red arrow in Figure 20c), and the second component was generated by TC JOANINHA to the east (black arrow in Figure 20c). In spite of the observed increase in wave height (and relatively constant wind field), a horizontal gradient in the aerosol concentration and size distribution at 100 m asl was not observed during this swell event. These results were predictable because the 1 m increase in wave height is relatively small and not expected to generate a significant difference in PMA emissions.




Figure 20. (a) Spectral density function of wave height simulated with WWW3 for legs A, B, C, and D. (b) Spectral density function of wave height from BOREAL UAS observations for legs A, B, C, and D. (c) Corresponding azimuth plot to wave elevation spectrum for leg A. The black (resp. red) arrow represents the wind direction (resp. current direction), and the scatter color are the wave crests. The length of the arrows has been scaled by 2.

Due to the low number of TCs passing in the immediate vicinity of Réunion Island during the TC season 2018–2019, the range of wind speeds and ocean wave conditions were not as large as expected to fully assess parameterizations for PMA emissions. Nevertheless, the datasets collected during this campaign have clearly demonstrated the scientific potential of the BOREAL UAS, and of its associated payload, to collect key observations in remote oceanic areas and in the vicinity to tropical cyclones.

3.2.2. GNSS Observations

A well-known (indirect) atmospheric application of GNSS measurement consists of measuring the atmospheric integrated water vapor (IWV) content from the delay induced by water vapor during the crossing of the earth's atmosphere by the GNSS signal [110]. Due to the fundamental role of water vapor in climate and weather dynamics, tropospheric GNSS measurements have rapidly become one of the main tools used by climatologists to monitor the evolution of the water vapor field at all spatio-temporal scales, but also to improve NWP model forecasts (see [111] for a recent review of current GNSS weather applications). The creation of the International GNSS Service (IGS) network in 1994 [112], which now includes more than 500 stations worldwide, has also enabled GNSS-derived measurements collected in all parts of the world to be widely disseminated to the scientific community. However, with only eight operational stations available prior to RNR-CYC (Figure 21), the density of public GNSS stations in the SWIO was the lowest of all TC basins.

During RNR-CYC, 10 new permanent stations were deployed, including one in the Seychelles (Aldabra), four in the Eparses Islands (three RNR-CYC stations at Juan de Nova, Tromelin and Europa and one shared station installed by OPGP in Glorieuses), and five in Madagascar (Diego-Suarez, Toamasina, Sainte-Marie, Nosy Be and Fort-Dauphin), with the goals of increasing the density of the GNSS network in the SWIO and to provide additional near-real-time IWV measurements at various locations in the basin (Figure 21). Although these stations have not yet been included into the IGS network (pending), most collected GNSS observations are transmitted at hourly time steps to IGN's data center, which already allows for routine real-time GNSS calculations and the wide dissemination of derived meteorological and geophysical products through IGN's permanent GNSS network (RGP, <ftp://rgpdata.ign.fr/pub/gnssmayotte>, accessed on 20 March 2021).

These new stations have already been shown to represent a fantastic asset for monitoring the spatio-temporal evolution of the water vapor field at local and regional scales [34,35], as well as to evaluate NWP forecasts [34]. A further modelling application is shown in Figure 22, which presents differences between GNSS-derived IWV observations and daily 6 h analyses of AROME-IO operational NWP systems [42] at 00:00, 06:00, 12:00, and 18:00 UTC at Aldabra (ALBR, The Seychelles), Antananarivo (ABPO, Madagascar), Réunion Island (RUN, France) and Sainte-Marie (MASM, Madagascar) during the year 2019. In order to evaluate the performance of the model at various time scales, 30-day (red), 10-day (yellow) and 3-day (black) moving averages were applied to both the model and GNSS data. GNSS observations were processed following the approach proposed by [35].



Figure 21. Map of public GNSS stations available in the SWIO as of March 2021 (top). Red squares show current IGS stations, green dots show GNSS stations installed during RNR-CYC, and blue triangles show stations to be installed in 2021 in the frame of the new project ESPOIRS (see Section 4). Pictures show GNSS stations installed in Fort-Dauphin (MAFD, upper-left), Tromelin (TRML, lower-left) and Aldabra (ALBR, lower-right). Sources: Google Earth (top); Olivier Bousquet (pictures)

Whatever the time scale, the bias error remained relatively constant at Sainte Marie (MASM, approximately -0.3 kg m^{-2}), Réunion (REUN, approximately 0.5 kg m^{-2}) and Antananarivo (ABPO, approximately -2 kg m^{-2}), but showed more variations in Aldabra (ALBR, between 0.6 and 1.14 kg m^{-2}). Model–observation differences did not show a clear seasonal dependency, except at Aldabra where maximum errors were observed in the middle of the winter, spring, and fall seasons. In the case of this atoll, consisting of land strips of 1–3 km width encircling a lagoon of nearly $30 \text{ km} \times 15 \text{ km}$, the land is considered as submerged by the model, which does not allow taking into account the effect of vegetation and of the diurnal cycle on the atmospheric moisture content. At Aldabra, atmospheric moisture changes in the model thus mostly depend on the variation of ocean temperatures. Interestingly, the sign of the bias error is location-dependent, suggesting that the model is not affected by systematic errors. The associated standard deviation errors also tends to decrease with the length of the smoothing period, because the latter minimized the impact of short-term variations. At short time scales (three days), one can

notice that the strongest discrepancies mostly occur during the TC (wet) season (days 1–120 and 330–365), in relation with the passage of tropical cyclones in the vicinity of the GNSS ground-based stations.




Figure 22. Time series of the differences between IWV contents observed by GNSS and AROME-IO analyses at Réunion Island (REUN, upper left), Sainte-Marie (MASM, upper right), Antananarivo (ABPO, lower left) and Aldabra (ALBR, lower right) throughout the year 2019. 30-day (red), 10-day (orange), and 3-day (black) moving averages were applied to IWV data to evaluate the impact of short and mid-term moisture variations on the model performance.

As already pointed out by [34], who investigated the seasonal variations of AROME-analyzed IWV errors at Diego-Suarez (DSUA), model–observation differences seem exacerbated for stations located in the vicinity of complex orography. Except for the particular case of Aldabra, the same behavior can be noticed here for the high-altitude stations of Réunion (600 m asl, REUN) and Antananarivo (2000 m asl, ABPO), which showed higher errors than for the flat island of Sainte-Marie (MASM). These errors might be related to the difficulties of the model to properly capture the modifications of air masses caused by the orography (lifting and subsiding motions) at the local scale due to an insufficient resolution of its topography. A possible way to correct for such errors could be to assimilate GNSS observations (zenithal delays) into the model. While the operational version of the AROME-IO NWP system does not allow for data assimilation, its research configuration includes a 3D-Var scheme that can be used to perform assimilation experiments [34]. Some studies are currently ongoing to determine whether the assimilation of observations collected from these new GNSS stations can help to reduce this moisture bias.

3.3. Spaceborne Observations

Visible and infrared satellite observations have long been the main source of knowledge for estimating some of the parameters (e.g., radius of maximum wind, various wind radii) characterizing the structure of tropical cyclones—these parameters are, in particular, inputs of the Dvorak method [113], used by TC forecast centers to produce best-track data and issue TC advisory and forecasts. Since the launch of the first Earth observation satellite in the early 1970s, the quality of ocean surface wind estimates in the vicinity of TCs has constantly been improved. This includes, for instance, the deployment of wind scatterometers [114,115] as well as of multifrequency radiometers [116,117] that both allow for a direct, and more precise, estimate of surface winds under TCs. More recently, a new generation of microwave radiometers has also been put into operation by ESA and the National Aeronautics and Space Administration (NASA) in the frame of the Soil Moisture

and Ocean Salinity (SMOS) and Soil Moisture Active Passive (SMAP) space missions, respectively. Thanks to their large coverage and revisit times, these new sensors, which allow for high-resolution (~40 km) surface wind speed sampling in all weather conditions, provide a unique description of TC structure during its whole lifetime [118], and have thus become a key source of information for TC forecasting centers.

In addition to these more or less conventional sensors, a new approach, based on the use of spaceborne synthetic aperture radars (SARs), is also being increasingly used to probe and quantify sea surface properties under extreme wind conditions. Designed in the mid-1990s, SAR systems were initially used for land applications [119], especially for monitoring earth deformation rates ranging from a few millimeters per year (e.g., for glaciers) up to 1 m per h (e.g., earthquakes and landslides). SAR potential for marine applications was already known since the launch of the first SAR satellite [120], but has significantly emerged with the launch of the Canadian Space Agency (CSA)'s RADARSAT-1 (1995) and ESA's Envisat (2002) satellites. Thanks to their unique capability to gather very high-resolution (up to a few meters) surface wind and roughness data in swaths of several hundred kilometers, spaceborne SARs have become key observing systems for monitoring, forecasting, and investigating the properties and evolution of TCs [120,121].

The deployment of a new generation of SAR systems equipped with polarization diversity on-board CSA's RADARSAT-2 (RS2, 2007) and ESA's Sentinel-1 (S1, 2014) satellites has allowed further improvements in the capabilities of these instruments to accurately map the variations of ocean surface winds in TC eyes and eyewalls [122,123]. Acquisitions made by ESA's satellites had, however, never been used to probe tropical cyclones until the implementation of the Satellite Hurricane Observations Campaign (SHOC, 2016–2017), in the Pacific and Atlantic basins. The intercomparison of aircraft reconnaissance wind data against S1 SAR measurements collected in CAT-5 hurricane IRMA (2017) during this experiment definitely demonstrated the capability of these instruments to thoroughly describe TC ocean boundary layer structures at high spatial resolution [124].

The extension of the SHOC program to the SWIO basin (referred to as SHOC-V2) was initiated in 2017, under the frame of RNR-CYC. SAR missions cannot continuously acquire wide swaths of data in high-resolution modes; therefore, a dedicated acquisition procedure was set up to collect data without impacting the operational duty cycle of the satellites. Acquisition requests were passed along to both MDA (MacDonald Dettwiler and Associates, the private company that owns the satellite RS2) and ESA S1 mission planner portals on a 24-to-48 h notice, based on satellite orbit and five-day track forecasts provided by RSMC La Réunion. In order to reduce the workload of participating space agencies, which contributed to this experimental program on a voluntary basis, acquisition requests were also generally limited to storms presenting a threat to SWIO populations.

Between 2017 and 2021, about 150 SAR images were acquired by S1A, S1B and RS2 satellites, allowing the sampling of 20 tropical storms and cyclones over four TC seasons (raw S1 data are available at <https://scihub.copernicus.eu>, accessed on 20 March 2021). Among these acquisitions, nearly 40% were directly obtained within the eye or the eyewall of the systems (Table 2, "hits"). SAR images were collected throughout the SWIO basin (Figure 23) at various storm evolution stages (cyclogenesis, intensification, dissipation). Collected wind data were used for nowcasting and best-track data reanalysis purposes at RSMC La Réunion, model verification [33] and assimilation in NWP systems [125] (this Special Issue).

Table 2. Number of Radarsat-2 and Sentinel-1 SAR acquisitions within the eyewall/eye (hits) of tropical storms (TSs) and tropical cyclones (TCs, ITC) that developed in the SWIO between February 2017 and 2021. * Acquisitions performed from 2021 onwards were made in the frame of the CYMS program (see Section 4).

Storm Name	Date	Hits	Storm Name	Date	Hits
Dineo (TC)	02/2017	2	Idai (ITC)	03/2019	3
Enawo (ITC)	03/2017	1	Joaninha (ITC)	03/2019	7
Ava (TC)	01/2018	3	Kenneth (ITC)	04/2019	1
Berguitta (ITC)	01/2018	4	Lorna (TC)	04/2019	4
Cebile (ITC)	01/2018	8	Belna (TC)	12/2019	5
Dumazile (ITC)	03/2018	5	Calvinia (TC)	12/2019	1
Eliakim (TS)	03/2018	1	Diane (TS)	01/2020	1
Alcide (ITC)	11/2018	1	Francisco (TS)	02/2020	1
Kenanga (ITC)	12/2018	2	Herold (ITC)	03/2020	1
Gelena (ITC)	02/2019	4	* Chalane (TS)	12/2021	3
Haleh (ITC)	03/2019	6	* Eloise (TC)	01/2021	2




Figure 23. Location of Sentinel-1 (A and B) and RADARSAT-2 swaths listed in Table 2.

Examples of SAR-derived surface wind fields obtained through the application of the retrieval algorithm proposed by [126] are presented hereafter. The first example was taken within TC JOANINHA, which developed in the northeastern part of the Mascarene Archipelago (Figure 17a) and reached its LMI in the vicinity of Rodrigues Island on 26 March 2019, with 10 min of maximum sustained winds of nearly 60 m s^{-1} (Figure 24a). The core structure of this system was observed seven times by the SAR systems deployed onboard satellites S1A/S1B and RS2 (Table 2) at various stages of its life cycle. Of particular interest is the image taken by RS2 on 28 March 2019 at 00:52 UTC (Figure 24b), when the system started to experience an eyewall replacement cycle (ERC). The latter, which often occurs when a TC reaches an intensity of 50 m s^{-1} , has long been recognized as one of the main mechanisms for intense TCs to intensify further [127,128]. During an ERC, outer rainbands strengthen and organize themselves into a ring of thunderstorms that progressively encircle the TC eyewall. The formation of this outer ring, usually referred to as the outer eyewall, eventually cuts off the supply of moisture and angular momentum that maintain convection around the eye, causing a weakening of the system and the eventual dissipation of the inner eyewall. The former ring is then replaced by the outer

ring, which gradually contracts and intensifies, often resulting in a more intense TC than prior to the ERC.

The analysis of the SAR-derived surface wind field shown in Figure 24b clearly indicates the presence of two concentric areas of maximum wind located both around the eye and in the east-to-southeastern quadrant of the system core. The inner structure (maximum wind speed of $\sim 50.2 \text{ m s}^{-1}$) corresponds to the eyewall of the storm, and the outer one (maximum wind speed of $\sim 45 \text{ m s}^{-1}$) to the cyclonic rainbands progressively wrapping around the eyewall that will eventually replace the inner wall. As mentioned previously, the completion of an ERC is often followed by a re-intensification of the system. However, in the present case, such re-intensification did not occur because TC JOANINHA rapidly encountered a strongly sheared environment that caused its dissipation a couple of days later.




Figure 24. Intensity of TC JOANINHA (color scale at bottom). (a) Wind speed evolution with respect to time as given by ATCF and (b) SAR wind retrieval (RS2) on 28 March 2020 at 00:52 UTC.

The second example is taken from TC IDAÏ (Figure 25), which is considered as the worst natural disaster that has ever affected Mozambique (as well as the surrounding countries of Zimbabwe and Malawi), and the deadliest storm ever recorded in the SWIO. This exceptional TC initiated as a tropical depression along the northern coast of Mozambique on 3 March 2019 (Figure 25a) that initially moved inland in the northwestern part of the country for a few days, with peak winds in the order of $10\text{--}15 \text{ m s}^{-1}$. On 7 March, it made a half-turn near the Mozambique–Malawi border and moved back towards the ocean. After entering the Mozambique Channel, on 9 March, the storm experienced a rapid intensification to reach intense TC intensity (10 min maximum sustained wind $>46 \text{ m s}^{-1}$), with winds gusts estimated at up to 70 m s^{-1} on 11 March. IDAÏ then reversed its track back for the second time in the immediate vicinity of the Eparses island of Juan de Nova

(located approximately 150 km off the west coast of Madagascar) and began its south-westward progression towards Mozambique. Shortly after this half-turn, it entered into a slight weakening trend following the beginning of its ERC. Right after its completion, IDAĭ immediately re-intensified and reached its LMI on 14 March, with a minimum central pressure of 940 hPa and (estimated) 10 min maximum sustained winds of $\sim 55 \text{ m s}^{-1}$. The storm then gradually weakened while progressing towards Beira, where it made landfall on 15 March at the TC stage.




Figure 25. As in Figure 24, but for (a) TC IDAĭ and (b) SAR images collected by S1A on 11 (02:43 UTC) and 14 (03:08 UTC) March 2019.

Numerous acquisitions were made in this system by S1 satellites, including three images collected directly in the core of the storm (Figure 25b). Two of these hits were made over the ocean by S1A, at the stage of intense TC (11 March at 02:43 UTC and 14 March at 03:08 UTC), while a third one was made at landfall by S1B on 14 March at 16:00 UTC (because 90% of this swath occurred over land, this third acquisition is not exploitable). SAR wind data shown in Figure 25 are the only high-resolution wind observations collected during the oceanic phase of this system—WMO’s JDN weather station broke down a week before the passage of the storm over the island and the weather station deployed at Beira in the frame of RNR-CYC was lost during landfall—therefore, these observations represent an invaluable asset for accurately assessing numerical model simulations of this storm [33].

SAR data collected in IDAĭ and other tropical cyclones can also meaningfully complement best-track (BT) data produced by RSMC La Réunion. According to RSMC La Réunion best-track, TC IDAĭ reached its LMI on 14 March 2019 at 00:00 UTC with maximum winds of nearly 55 m s^{-1} . This value is in good agreement with the maximum intensity measured by S1B a few hours later ($\sim 52 \text{ m s}^{-1}$). SAR data, however, show that these strong wind values were only observed in the eastern quadrant of the storm’s core, but that the average wind speed throughout the eyewall was significantly less intense and mostly comprised between 42 and 45 m s^{-1} . On 11 March, the agreement between best-track and SAR-derived

wind data was less good, with maximum wind speed in the order of 50 m s^{-1} for the BT and 60 m s^{-1} for SAR observations. According to SAR observations, the strongest wind speed values were also more or less uniformly distributed throughout the eyewall. Hence, the comparison of the two SAR images suggests that the destructive potential, and overall global intensity of the system, was probably much greater on 11 March, despite a 10% lower maximum wind speed (50 m s^{-1} vs. 55 m s^{-1}) in BT data.

4. Conclusions and Perspectives

ReNovRisk-Cyclone (RNR-CYC) is an ambitious international research program aimed at developing regional cooperation in the SWIO with emphasis on the observation and modeling of tropical cyclones at both current and future time horizons. The observation component of RNR-CYC, presented in detail in this paper, has enabled the collection of numerous innovative oceanic (gliders, sea-turtle borne and seismometer-derived data) and atmospheric (UAS and SAR-derived wind data) measurements, together with more conventional observations (GNSS-derived IWV, atmospheric radiosoundings, ADCP, and wave gauge swell measurements) to investigate tropical cyclones and their environment from the local to the basin scale. Its mesoscale modelling and climate components, presented in more detail in the companion paper [33], also allowed for the development of innovative modelling systems, while its outreach component significantly increased regional collaboration between SWIO countries affected by TC hazards. The innovative data collected in the SWIO during RNR-CYC have also proven particularly useful to evaluate the performance of both research and operational TC forecasting models in this under-sampled region of the world. SAR measurements, which allow for accurate evaluations of TC forecasts, can also be assimilated into NWP systems [126]; in situ and remote swell measurements can be relied upon to evaluate wave model performance; while biologging measurements are priceless to assess ocean model representation of the ocean mixed layer in cyclonic conditions.

The promising results obtained during the 3.5-year observation component of RNR-CYC have already led to the development of numerous new research projects aimed at maintaining some of these novel measurements and further reinforcing overall observation capabilities in the SWIO beyond the end of this program (July 2021). Among the principal new research projects directly arising from RNR-CYC, one can mention STORM-IO (Sea Turtle for Ocean Research and Monitoring in the Indian Ocean), ESPOIRS (Etude des systèmes précipitants de l'océan Indien par radar et satellites—Studies of Indian Ocean precipitation systems by radars and satellites), and MAP-IO (Marion Dufresne Atmospheric Program—Indian Ocean).

STORM-IO (starting May 2021) will enable the extension ST-borne measurements conducted from Réunion Island to the whole Indian Ocean (Figure 26), in collaboration with the Terres Australes et Antarctiques Françaises (TAAF) administration (in Eparses Islands of Juan de Nova and Tromelin), Kelonia (Réunion Island), and five marine reservations in Comoros (Moheli), Madagascar (Nosy Tanikely), Seychelles (Aldabra), Mozambique (Ponto di Ouro), and South Africa (iSimanliso). This transdisciplinary project, constructed in close cooperation with ST specialists and oceanographers, will extend the research initiated during RNR-CYC to: (i) investigate Indian Ocean properties and spatio-temporal variability; and (ii) improve knowledge of the ecology of the five species of sea turtles living in the Indian Ocean. A particular emphasis will be put on the observation of mesoscale eddies and coastal currents that develop in the Mozambique Channel (to assess their impact on water mass distribution, transport, and mixing, as well as their role in the dynamics of the Greater Agulhas current system), and on the assimilation of collected ST-borne data in global and limited-area configurations of the ocean model NEMO [129]. This new component of the global STORM program, which is now strongly supported by the EU, UNESCO (through the GOOS/ AniBOS), CNES and CNRS, among others, will be conducted for the many years to come, and is likely to be extended to other ocean basins in the mid-term.




Figure 26. Release locations and sea turtle species to be equipped in the frame of the STORM-IO project: Ei: *Eretmochelys imbricata* (Hawksbill); Cm: *Chelonia mydas* (Green); Cc: *Caretta caretta* (Loggerhead); Lo: *Lepidochelys olivacea* (Olive Ridley); Dc: *Dermochelys coriacea* (Leatherback). Sources: background, Google Earth; Pictures, Kelonia

ESPOIRS (started in December 2020) will carry on with the densification of the GNSS water vapor observation network initiated in the framework of RNR-CYC's sub-program IOGA⁴MET through the deployment of five additional stations in Madagascar and Mozambique (Figure 21). Some existing, or soon to be deployed, GNSS stations will also be upgraded with co-located oceanographic sensors to monitor sea level on a regional scale. In the mid-term, these instruments will be integrated into the Global Sea Level Observing System (GLOSS, <https://www.gloss-sealevel.org/>, accessed on 20 March 2021) network, which monitors sea level rise on a global scale, and into the IGS network, which disseminates GNSS data to the worldwide scientific community. ESPOIRS also includes an ambitious local component aimed at collecting wind and precipitation measurements in tropical cyclones with a transportable polarimetric weather radar to be deployed in Réunion Island (2021), the Seychelles (Mahe, 2022), and Madagascar (Tamatave or Diego Suarez, 2023). This weather radar component will notably focus on the impact of orography on tropical cyclones (or outer cyclonic rainbands in the Seychelles) [130], and will help to develop regional cooperation in the field of atmospheric remote sensing.

The goal of the MAP-IO program (started in January 2021) is to deploy an atmospheric and marine biology observatory in the Indian and Southern Oceans onboard the French vessel Marion Dufresne (Figure 27), operated by the TAAF administration (for its logistical needs in the Eparses Island and French Austral Territories) and IFREMER (for scientific campaigns in the IO). With the deployment of nearly 20 new sensors on-board this RV (e.g., cytometer, titrator, thermosalinograph, cytometer, titrator, NO_x, CO, CO₂, O₃ and CH₄ analyzers, GNSS, UV radiometer, CCN, photometer, weather station), observations at the OA interface and within the atmospheric column will be collected over the long term in this region particularly sensitive to the impact of climate change (note that all observations will be transmitted in near-real-time through the RV onboard satellite communication system). The data collected in the Southern Ocean for sea and wind conditions close to those encountered in tropical cyclones will also be used to carry on with the evaluation of OA exchange parameterizations (turbulent fluxes, aerosol, and sea-spray emissions) in extreme conditions [33]. This exploratory research project, which benefits from strong

support from the EU, IFREMER, CNRS, TAAF and Réunion University, among others, is likely to be maintained in the long term.



Figure 27. Picture of the RV Marion Dufresne anchored at Mayotte during TAAF's Eparses Island rotation in April 2019. Source: Olivier Bousquet

Finally, one can also mention ESA's pre-operational program CYclone Monitoring Service based on Sentinel-1 (CYMS, <https://www.esa-cyms.org>, accessed on 20 March 2021), which is aiming to extend SAR acquisitions to all TC basins and provide real-time ocean surface wind measurements to further demonstrate SAR potential for operational TC forecasting. This new program, motivated in large part by the results obtained in RNR-CYC, will further develop understanding of the impact of TCs on the earth system, and likely open new scientific applications of SAR measurements.

Author Contributions: Conceptualization O.B. and G.B.; software, C.B., S.B., P.T., J.P., J.C. and F.C.; data acquisition and processing, O.B., G.B., E.C., E.R., R.C., G.R., J.D., A.M. (Alexis Mouche), R.H., S.C., E.L., D.M. and N.M.; writing—original draft preparation, O.B., G.B., E.C., R.C., E.R., G.R. and C.B.; writing—review and editing, all co-authors; supervision, O.B.; project administration, O.B.; funding acquisition, O.B., P.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research was principally funded by the European Union, the Regional Council of Réunion Island, and the French State through INTERREG-V Océan Indien 2014–2020 projects “ReNovRisk-Cyclones and Climate Change” (TF) and “ReNovRisk Cyclones and Precipitation” (TF), as well as by Université de La Réunion and Centre National de la Recherche Scientifique (CNRS) in the frame of the “Contrat Plan Etat Région” (CPER) “ReNovRisk”. The gliders were provided, operated, and co-funded by CNRS DT-INSU (Direction Technique—Institut National des Sciences de l'Univers). The seismic stations deployed in the “Rivière des Pluies” network (code ZF) were provided by the INSU-RESIF/SISMOB instrumental pool. The provision and operating of the BOREAL UAS were co-funded by Météo-France. The installation of GNSS stations on Eparses islands was funded by the Eparses consortium of the Terres Australes et Antarctiques Françaises (TAAF) administration under the frame of project “IOGA4MET-EI”. Part of the STORM operations were also co-funded by CNRS-INSU under the frame of the LEFE project “PreSTORM”. Access to RADARSAT-2 data was supported by Institut Mines Télécom and by public funds (Ministère de l'Education Nationale, de l'Enseignement Supérieur et de la Recherche, FEDER, Région Bretagne, Conseil Général du Finistère, Brest Métropole) received in the framework of the VIGISAT project managed by Groupement Bretagne Télédétection (GIS BreTel – Brittany Remote Sensing).

Institutional Review Board Statement: All sea turtles equipped in this programme were handled by qualified and officially accredited personnel from Kelonia's care center. The used tags meet all

the requirements of the international conventions for the protection of sea turtles and were directly purchased from manufacturers specializing in marine biology and biologging.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Some of the data used in this study are available in public repositories: <https://scihub.copernicus.eu/> and <https://cycloblogs.ifremer.fr/> (S1 and RS2 SAR data); http://www.meteo.fr/temps/domtom/La_Reunion/webcmrs9.0/anglais/index.htm (best-track data); <ftp://rgpdata.ign.fr/pub/gnssmayotte> (GNSS data); <http://seismology.resif.fr/> (seismic data—FDSN network codes PF and ZF), accessed on 20 March 2021. Other data used in this study are not yet publicly available due to temporary use restrictions by data owners (these data, available on request from the corresponding author, will be soon deposited in the project’s data repository).

Acknowledgments: The authors are grateful to the numerous students, researchers, and engineers who participated in the field phase of the project, as well as to the staff of IGN, for processing GNSS data, Kelonia and Centre d’Etude et de Découverte des Tortues Marines (CEDTM), for equipping sea turtles, CNRS (DT-INSU), for operating the gliders, and Centre National de Recherches Météorologiques (CNRM, Météo-France), for operating the BOREAL UAS. The BOREAL UAS had been developed in the MIRIAD project (Système de Mesures scientifiques de flux de surface en milieu maritime embarqué sur Drone), financed by the European Union and the Occitanie Region. The SHOC initiative has been possible thanks to SAR data access supported by the ESA Sentinel-1 mission ground segment team and GIS BreTel. Sentinel-1 is part of the European space component of Copernicus European program. RADARSAT-2 is a commercial mission and data were provided by MDA’s Geospatial Services (<https://mdacorporation.com/geospatial/international> (accessed on 18 April 2021)). We are grateful to the OVPF (Observatoire Volcanologique du Piton de la Fournaise) and to IGP colleagues who maintain the PF seismic stations and to F.R. Fontaine, E. Delcher, and A. Gonzalez for the Riviere des Pluies seismic data acquisition (<http://dx.doi.org/10.15778/RESIF.ZF2015>, accessed on 20 March 2021). We would also like to warmly thank Mayeul Dalleau (CEDM) for his important contribution in the design of the STORM sub-programme, and Navalona Ravoson (Université d’Antananarivo), for contributing to seismic data analysis. Finally, we would also like to express our warmest thanks to Mrs Sandrine Prunier and Nina Breznik, who successfully ensured the administrative management of this project.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Jiang, H.; Oyama, R.; Bousquet, O.; Vigh, J.; Huang, Y.-H.; Smith, R.; Corbosiero, K.L.; Hendricks, E.; Kepert, J.D.; Miyamoto, Y.; et al. A Review of Tropical Cyclone Intensity Change (2014–2018). Part II: Rapid Intensification via Internal Influences. *Atmosphere* **2021**. Submitted.
- Vigh, J.L.; Huang, Y.-H.; Miyamoto, Y.; Li, Q. A Review of Intensity Change (2014–2018). Part III: Internal Influences. *J. Meteorol. Soc. Jan.* **2021**. Submitted.
- Courtney, J.B.; Langlade, S.; Barlow, S.; Birchard, T.; Knaff, J.A.; Kotal, S.; Kriat, T.; Lee, W.; Pasch, R.; Sampson, C.R.; et al. Operational perspectives on tropical cyclone intensity change Part 2: Forecasts by operational agencies. *Trop. Cyclone Res. Rev.* **2019**, *8*, 226–239. [[CrossRef](#)]
- Heming, J.; Prates, F.; Bender, M.A.; Bowyer, R.; Cangialosi, J.; Caroff, P.; Coleman, T.; Doyle, D.J.; Dube, A.; Faure, G.; et al. Review of recent progress in tropical cyclone track forecasting and expression of uncertainties. *Trop. Cyclone Res. Rev.* **2019**, *8*, 181–218. [[CrossRef](#)]
- Yablonsky, R.M. Ocean Component of the HWRF Coupled Model and Model Evaluation. In *Advanced Numerical Modeling and Data Assimilation Techniques for Tropical Cyclone Prediction*; Mohanty, U.C., Gopalakrishnan, S.G., Eds.; Springer: Dordrecht, The Netherlands, 2016. [[CrossRef](#)]
- Mogensen, K.S.; Magnusson, L.; Bidlot, J.-R. Tropical cyclone sensitivity to ocean coupling in the ECMWF coupled model. *J. Geophys. Res. Oceans* **2017**, *122*, 4392–4412. [[CrossRef](#)]
- Feng, X.; Klingaman, N.P.; Hodges, K.I. The effect of atmosphere–ocean coupling on the prediction of 2016 western North Pacific tropical cyclones. *Q. J. R. Meteorol. Soc.* **2019**, *145*, 2425–2444. [[CrossRef](#)]
- Vellinga, M.; Copsey, D.; Graham, T.; Milton, S.; Johns, T. Evaluating Benefits of Two-Way Ocean–Atmosphere Coupling for Global NWP Forecasts. *Weather Forecast.* **2020**, *35*, 2127–2144. [[CrossRef](#)]
- Bousquet, O.; Dalleau, M.; Bocquet, M.; Gaspar, P.; Bielli, S.; Ciccione, S.; Remy, E.; Vidard, A. Sea Turtles for Ocean Research and Monitoring: Overview and Initial Results of the STORM Project in the Southwest Indian Ocean. *Front. Mar. Sci.* **2020**, *7*, 859. [[CrossRef](#)]

10. Leroux, M.-D.; Wood, K.; Elsberry, R.L.; Cayan, E.O.; Hendricks, E.; Kucas, M.; Otto, P.; Rogers, R.; Sampson, B.; Yu, Z. Recent Advances in Research and Forecasting of Tropical Cyclone Track, Intensity, and Structure at Landfall. *Trop. Cyclone Res. Rev.* **2019**, *7*, 85–105. [[CrossRef](#)]
11. Pianezze, J.; Barthe, C.; Bielli, S.; Tulet, P.; Jullien, S.; Cambon, G.; Bousquet, O.; Claeys, M.; Cordier, E. A New Coupled Ocean-Waves-Atmosphere Model Designed for Tropical Storm Studies: Example of Tropical Cyclone Bejisa (2013–2014) in the South-West Indian Ocean. *J. Adv. Model. Earth Syst.* **2018**, *10*, 801–825. [[CrossRef](#)]
12. Pant, V.; Prakash, K.R. Response of Air–Sea Fluxes and Oceanic Features to the Coupling of Ocean–Atmosphere–Wave during the Passage of a Tropical Cyclone. *Pure Appl. Geophys. Pageoph.* **2020**, *177*, 3999–4023. [[CrossRef](#)]
13. Prakash, K.R.; Pant, V.; Nigam, T. Effects of the sea surface roughness and sea spray-induced flux parameterization on the simulations of a tropical cyclone. *J. Geophys. Res. Atmos.* **2019**, *124*, 14037–14058. [[CrossRef](#)]
14. Mavume, A.F.; Rydberg, L.; Lutjeharms, J.R.E. Climatology of tropical cyclones in the South-West Indian Ocean; landfall in Mozambique and Madagascar. *West. Indian Ocean J. Mar. Sci.* **2008**, *8*, 15–36.
15. Matyas, C.J. Formation and movement of tropical cyclones in the Mozambique channel. *Int. J. Clim.* **2014**, *35*, 375–390. [[CrossRef](#)]
16. Leroux, M.-D.; Meister, J.; Mekies, D.; Dorla, A.-L.; Caroff, P. A Climatology of Southwest Indian Ocean Tropical Systems: Their Number, Tracks, Impacts, Sizes, Empirical Maximum Potential Intensity, and Intensity Changes. *J. Appl. Meteorol. Clim.* **2018**, *57*, 1021–1041. [[CrossRef](#)]
17. Vialard, J.; Foltz, G.R.; McPhaden, M.J.; Duvel, J.-P.; Montégut, C.D.B. Strong Indian Ocean sea surface temperature signals associated with the Madden-Julian Oscillation in late 2007 and early 2008. *Geophys. Res. Lett.* **2008**, *35*, L19608. [[CrossRef](#)]
18. Hermes, J.C.; Reason, C.J.C. The sensitivity of the Seychelles–Chagos thermocline ridge to large-scale wind anomalies. *ICES J. Mar. Sci.* **2009**, *66*, 1455–1466. [[CrossRef](#)]
19. Yokoi, T.; Tozuka, T.; Yamagata, T. Seasonal Variation of the Seychelles Dome. *J. Clim.* **2008**, *21*, 3740–3754. [[CrossRef](#)]
20. Mawren, D.; Hermes, J.; Reason, C.J.C. Exceptional Tropical Cyclone Kenneth in the Far Northern Mozambique Channel and Ocean Eddy Influences. *Geophys. Res. Lett.* **2020**, *47*, e2020GL088715. [[CrossRef](#)]
21. Emerton, R.; Cloke, H.; Ficchi, A.; Hawker, L.; de Wit, S.; Speight, L.; Prudhomme, C.; Rundell, P.; West, R.; Neal, J.; et al. Emergency flood bulletins for Cyclones Idai and Kenneth: A critical evaluation of the use of global flood forecasts for international humanitarian preparedness and response. *Int. J. Disaster Risk Reduct.* **2020**, *50*, 101811. [[CrossRef](#)]
22. Matera, M. World Bank’s Cyclone IDAI & Kenneth Emergency Recovery and Resilience Project 171040. 2019. Available online: <http://documents1.worldbank.org/curated/en/727131568020768626/pdf/Project-Information-Document-Mozambique-Cyclone-Idai-Kenneth-Emergency-Recovery-and-Resilience-Project-P171040.pdf> (accessed on 21 April 2021).
23. Tulet, P.; Aunay, B.; Barruol, G.; Barthe, C.; Belon, R.; Bielli, S.; Bonnardot, F.; Bousquet, O.; Cammas, J.-P.; Cattiaux, J.; et al. ReNovRisk: A multidisciplinary programme to study the cyclonic risks in the South-West Indian Ocean. *Nat. Hazards* **2021**, 1–33. [[CrossRef](#)]
24. Sharmila, S.; Walsh, K.J.E. Recent poleward shift of tropical cyclone formation linked to Hadley cell expansion. *Nat. Clim. Chang.* **2018**, *8*, 730–736. [[CrossRef](#)]
25. Camargo, S.J.; Wing, A.A. Increased tropical cyclone risk to coasts. *Science* **2021**, *371*, 458–459. [[CrossRef](#)]
26. Seidel, D.J.; Fu, Q.; Randel, W.J.; Reichler, T.J. Widening of the tropical belt in a changing climate. *Nat. Geosci.* **2008**, *1*, 21–24. [[CrossRef](#)]
27. Yang, H.; Lohmann, G.; Lu, J.; Gowan, E.J.; Shi, X.; Liu, J.; Wang, Q. Tropical Expansion Driven by Poleward Advancing Midlatitude Meridional Temperature Gradients. *J. Geophys. Res. Atmos.* **2020**, *125*, 033158. [[CrossRef](#)]
28. Kossin, J.P.; Emanuel, K.A.; Vecchi, G.A. The poleward migration of the location of tropical cyclone maximum intensity. *Nat. Cell Biol.* **2014**, *509*, 349–352. [[CrossRef](#)]
29. Kossin, J.P.; Emanuel, K.A.; Camargo, S.J. Past and Projected Changes in Western North Pacific Tropical Cyclone Exposure. *J. Clim.* **2016**, *29*, 5725–5739. [[CrossRef](#)]
30. Sun, J.; Wang, D.; Hu, X.; Ling, Z.; Wang, L. Ongoing Poleward Migration of Tropical Cyclone Occurrence Over the Western North Pacific Ocean. *Geophys. Res. Lett.* **2019**, *46*, 9110–9117. [[CrossRef](#)]
31. Krupar, R.J.; Smith, D.J. Poleward Migration of Tropical Cyclone Activity in the Southern Hemisphere: Perspectives and Challenges for the Built Environment in Australia. In *Hurricane Risk*; Collins, J., Walsh, K., Eds.; Springer: Cham, Switzerland, 2019; Volume 1. [[CrossRef](#)]
32. Cattiaux, J.; Chauvin, F.; Bousquet, O.; Malardel, S.; Tsai, C.-L. Projected Changes in the Southern Indian Ocean Cyclone Activity Assessed from High-Resolution Experiments and CMIP5 Models. *J. Clim.* **2020**, *33*, 4975–4991. [[CrossRef](#)]
33. Barthe, C.; Bousquet, O.; Bielli, S.; Tulet, P.; Pianezze, J.; Claeys, M.; Tsai, C.-L.; Thompson, C.; Bonnardot, F.; Chauvin, F.; et al. Impact of tropical cyclones on inhabited areas of the SWIO basin at present and future horizons. Part 2: Modelling component of the research program RENOVIRISK-CYCLONE. *Atmosphere* **2021**. submitted to this Special Issue.
34. Bousquet, O.; Lees, E.; Durand, J.; Peltier, A.; Duret, A.; Mekies, D.; Boissier, P.; Donal, T.; Fleischer-Dogley, F.; Zakariasy, L. Densification of the Ground-Based GNSS Observation Network in the South-west Indian Ocean: Current Status, Perspectives, and Examples of Applications in Meteorology and Geodesy. *Front. Earth Sci.* **2020**, *8*, 609757. [[CrossRef](#)]
35. Lees, E.; Bousquet, O.; Roy, D.; De Bellevue, J.L. Analysis of diurnal to seasonal variability of Integrated Water Vapour in the South Indian Ocean basin using ground-based GNSS and fifth-generation ECMWF reanalysis (ERA5) data. *Q. J. R. Meteorol. Soc.* **2021**, *147*, 229–248. [[CrossRef](#)]



atmosphere



Article

Cloud Radar Observations of Diurnal and Seasonal Cloudiness over Reunion Island

Jonathan Durand, Edouard Lees, Olivier Bousquet, Julien Delanoë and François Bonnardot



<https://doi.org/10.3390/atmos12070868>

Article

Cloud Radar Observations of Diurnal and Seasonal Cloudiness over Reunion Island

Jonathan Durand ¹, Edouard Lees ¹, Olivier Bousquet ^{1,2,*}, Julien Delanoë ³ and François Bonnardot ⁴

¹ Laboratoire de l'Atmosphère et des Cyclones (UMR8105 LACy), Université de La Réunion, CNRS, Météo-France, 97400 Saint-Denis, France; jonathan.durand@univ-reunion.fr (J.D.); edouard.lees@meteo.fr (E.L.)

² Institute for Coastal Marine Research (CMR), Nelson Mandela University, Port-Elizabeth 6001, South Africa

³ Laboratoire Atmosphère, Milieux et Observations Spatiales (UMR 8190 LATMOS), CNRS/Sorbonne Université/USVQ, 78280 Guyancourt, France; Julien.Delanoë@latmos.ipsl.fr

⁴ Direction Interrégionale de Météo-France pour l'Océan Indien, Saint-Denis, 97490 Sainte-Clotilde, France; francois.bonnardot@meteo.fr

* Correspondence: olivier.bousquet@meteo.fr

Abstract: In November 2016, a 95 GHz cloud radar was permanently deployed in Reunion Island to investigate the vertical distribution of tropical clouds and monitor the temporal variability of cloudiness in the frame of the pan-European research infrastructure Aerosol, Clouds and Trace gases Research InfraStructure (ACTRIS). In the present study, reflectivity observations collected during the two first years of operation (2016–2018) of this vertically pointing cloud radar are relied upon to investigate the diurnal and seasonal cycle of cloudiness in the northern part of this island. During the wet season (December–March), cloudiness is particularly pronounced between 1–3 km above sea level (with a frequency of cloud occurrence of 45% between 12:00–19:00 LST) and 8–12 km (with a frequency of cloud occurrence of 15% between 14:00–19:00 LST). During the dry season (June–September), this bimodal vertical mode is no longer observed and the vertical cloud extension is essentially limited to a height of 3 km due to both the drop-in humidity resulting from the northward migration of the ITCZ and the capping effect of the trade winds inversion. The frequency of cloud occurrence is at its maximum between 13:00–18:00 LST, with a probability of 35% at 15 LST near an altitude of 2 km. The analysis of global navigation satellite system (GNSS)-derived weather data also shows that the diurnal cycle of low- (1–3 km) and mid-to-high level (5–10 km) clouds is strongly correlated with the diurnal evolution of tropospheric humidity, suggesting that additional moisture is advected towards the island by the sea breeze regime. The detailed analysis of cloudiness observations collected during the four seasons sampled in 2017 and 2018 also shows substantial differences between the two years, possibly associated with a strong positive Indian Ocean Southern Dipole (IOSD) event extending throughout the year 2017.

Keywords: cloud radar; BASTA; Indian Ocean Subtropical Dipole; Reunion Island; integrated water vapor; Southwest Indian Ocean; GNSS; cloudiness variability; tropical clouds



Citation: Durand, J.; Lees, E.; Bousquet, O.; Delanoë, J.; Bonnardot, F. Cloud Radar Observations of Diurnal and Seasonal Cloudiness over Reunion Island. *Atmosphere* **2021**, *12*, 868. <https://doi.org/10.3390/atmos12070868>

Academic Editors: Lawrence D. Carey and Filomena Romano

Received: 29 May 2021

Accepted: 29 June 2021

Published: 3 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Because cloudiness plays a major role in distributing the radiative energy available in the atmosphere [1,2], knowledge of cloud properties is crucial for anticipating the impacts of cloud cover and, in particular, low-level cloud feedbacks in climate change projections [3]. A better knowledge of the vertical distribution of clouds is also important for numerical modelling, as errors in the location and vertical structure of clouds, which directly impact the radiative balance, are considered among the main sources of uncertainty in climate and weather numerical weather prediction (NWP) models [4–6].

In the tropics, cloudiness is principally determined by the space–time distribution of the lower-tropospheric moisture, which controls the total water content and buoyancy of

atmospheric parcels. Humidity is essentially driven by the seasonal migration of the Inter-Tropical Convergence Zone (ITCZ), and its associated circulation shifts, as well as by large scale anomalies that influence atmospheric circulation at both intra-seasonal (Madden–Julian Oscillation [7,8]), and interannual (e.g., El Niño Southern Oscillation, Indian Ocean Dipole, Indian Ocean Subtropical Dipole) timescales.

Thanks to their ability to cover vast areas of the globe, spaceborne cloud radars (e.g., CloudSat) have significantly improved our understanding of the spatio-temporal distribution of clouds throughout the planet and are widely considered as one of the most efficient tools to study cloud properties at the global scale [9]. These space-borne observations are nevertheless subject to some limitations, especially at low-level, due to ground (or sea) clutter echoes [10–12] and difficulties to penetrate thick mid-level clouds to observe the low cloud beneath (due to their short wavelength). In this regard, ground-based cloud radars, which feature higher spatio-temporal time resolution than space-borne sensors, are essential to complete (and evaluate) space-borne observations, especially at the local scale [13]. Relatively few studies have however been conducted so far to investigate tropical cloud properties from ground-based cloud radars. Moreover, if one except a few studies conducted from the Atmospheric Radiation Measurement (ARM) Tropical Western Pacific (TWP) sites [14–16], the latter were essentially conducted in the northern hemisphere due to the overall lack of observation facilities available south of the equator. This includes measurements conducted in Niger [17], in the Indian Western Ghats [18], or from the Barbados Cloud Observatory (BCO [19]) to investigate the seasonality and vertical distribution of low-level clouds [20,21]. This lack of observations was further exacerbated in the last few years following the dismantling, in 2013 (Nauru) and 2014 (Darwin and Manus) of the three permanent Pacific ARM TWP sites [22]. As a consequence, the Atmospheric Physics Observatory (OPAR) of Reunion Island, a small (2512 km²) volcanic French overseas territory located in the Southwest Indian Ocean (SWIO), has now become the principal atmospheric research facility in the southern hemisphere.

Due to its location, located both at the edge of the subtropical jet and in the subsidence region of the southern hemisphere Hadley cell, Reunion Island (20.8° S, 55.5° E) is particularly well suited to study tropical and subtropical influences on the general circulation. For this reason, OPAR's instrumentation is principally designed to observe high troposphere—low stratosphere interactions from a unique ensemble of lidars and radiometric sensors deployed at the high-altitude Maïdo observatory [23]. In order to extend cloud and precipitation observation capabilities, a new coastal experimental site was, however, recently integrated to OPAR in the frame of the pan-European research infrastructure “Aerosol, Clouds and Trace gases Research InfraStructure” (ACTRIS, <https://www.actris.eu>, accessed on 1 July 2021), a programme aiming at collecting high-quality observation data to constrain predictive models and improve global forecasts of the atmosphere. This new observation facility, located on the northern coast of the island at Saint-Denis de La Réunion (Figure 1), was notably equipped, in 2016, with a 95 GHz radar BASTA (Bistatic rAdar SysTem for Atmospheric Studies; [24]) to collect cloud observations at high spatial and temporal resolutions. This new site is, in particular, well-adapted to investigate coastal tropical clouds, which are known to occur over different mid-tropospheric conditions than over the open ocean and continental areas due to the combined effects of land–sea breeze interactions [25].

In this study, reflectivity observations collected during the two first years of operation of the cloud radar BASTA are used to investigate the characteristics of cloudiness in Reunion Island as well as its relationship with local and regional tropospheric humidity patterns inferred from radiosounding, Global Navigation Satellite System (GNSS) and high-resolution reanalysis data. Although limited in time, the 2-y cloud radar dataset used in this study allows us to investigate, for the first time in the SWIO, the vertical cloud properties over a remote and climatologically important location of the Southern Hemisphere, which was identified as a key component in the climate system by ACTRIS research infrastructure.




Figure 1. The cloud radar BASTA-REUNION located in Saint-Denis de La Réunion on the rooftop of Reunion University (55.48° E, 20.9° S).

This paper is organized as follows: information about data used in this study are given in Section 2; the distribution and analysis of the seasonal cloud cover over Saint-Denis between November 2016 and October 2018 are presented in Section 3; possible relationships between observed cloud patterns and large-scale environment are thus investigated in Section 4, while Section 5 presents our conclusions and perspectives.

2. Data and Methods

2.1. Cloud Radar Observations

The present study is principally based on continuous observations collected between November 2016 and October 2018 by a cloud radar BASTA. BASTA is a 95 GHz vertically pointing bistatic Doppler radar (distinct antennas for transmission and reception) developed by LATMOS (“Laboratoire Atmosphère, Milieux et Observations Spatiales”) for cloud and fog studies [24,26]. A dedicated version of this radar (referred to as BASTA-REUNION) was deployed in Reunion Island in November 2016 at the main campus facility of Reunion University, located in Saint-Denis approximately 100 m above ground level (Figure 1). BASTA-REUNION provides times series of vertical reflectivity and Doppler velocity profiles up to an altitude of 24 km. Reflectivity profiles consist of a combination of data collected in four distinct operating modes that are each activated over a period of integration of 3 s and later recombined to reconstruct the cloud structure throughout the

tropospheric layer every 12 s. The first operating mode, dedicated to fog and low cloud observations, runs at the resolution of 12.5 m and provides information up to an altitude of 12 km. The second mode, which features a vertical resolution of 25 m, is used to investigate mid-level clouds between 6 km and 18 km. Finally, the last two modes (100 m vertical resolution) are used to sample high-level clouds up to 18 km and 24 km, respectively—the latter mode, which is slightly more sensitive, was specifically implemented for tropical regions. Nearly four million reconstructed vertical reflectivity profiles are used in this study, which represent ~80% of the total number of profiles theoretically expected over a 2-year period.

Due to its operating frequency, BASTA-REUNION can be strongly affected by radar attenuation resulting from the absorption or reflection of transmitted signals in precipitation areas. This phenomenon, which occurs at most radar frequencies, reflects the fact that some of the energy is often lost to scattering and absorption when the radar pulse penetrates in heavy precipitation areas. At 95 GHz, the frequency of BASTA-REUNION, uncertainties can be introduced in the observed cloud properties and occasionally can result in the extinction of the radar signal in a heavy precipitation regime. As will be seen later, this phenomenon nevertheless remains relatively uncommon in the Saint-Denis area and does not significantly affect the statistical analysis of cloud radar data that will be presented hereafter. We thus chose to not discriminate between precipitating and non-precipitating clouds so as to be able to also identify the occurrence of rainfall in radar observations—as fog and low-level stratus never occur over Saint-Denis, radar data collected below 1 km always reflect the presence of precipitation.

During the austral summer, the presence of the Intertropical Convergence Zone (ITCZ) nearby Reunion Island induces low-level convergence of highly buoyant air that favors the development of deep convection. The interactions of these unstable air masses with the complex orography of the island can generate very intense precipitation due, in particular, to the passage of tropical cyclones nearby [27]. In winter, as the ITCZ moves back northwards, the local weather becomes mostly influenced by the subsided southern branch of the Hadley cell. The resulting temperature inversion tends to inhibit atmospheric upward vertical motion above ~3 km [28,29] and is further amplified by the trade wind inversion [30–32]. Due to its complex orography, Reunion Island is also affected all year long by thermally induced sea breeze circulations that can force onshore moist flow at low levels and further favors the development of low-level clouds during daytime [30,33]. Examples of observations collected by BASTA-REUNION in Austral summer (26 March 2017), Austral winter (1 August 2017) as well as during cyclonic conditions (4–6 March 2018) are presented hereafter to illustrate the general cloudiness conditions associated with the various large-scale circulation patterns that drive cloud formation in Reunion Island.

In Austral summer (Figure 2a, 26 March 2017), radar observations show the existence of a strong diurnal cycle characterized by low-level clouds forming early in the morning and eventually dissipating in the late evening. Convective clouds, resulting from the high convective instability that prevails over the tropical SWIO basin during summer, develop in the early afternoon and extend up to the altitude of 10 km (all heights are given above mean sea level). Precipitation, as evidenced by associated time series of collocated rain gauge measurements, is maximised around mid-day (9–11 UTC, 13–15 LST) and dissipates in the early evening. During winter (Figure 2b, 1 August 2017), low-level clouds also form in the early morning (near 03:00 UTC/07:00 LST), but never evolve into deep convective clouds due to the strong capping effect. These low-level clouds, which consist of small, non-precipitating, scattered cumulus, are not advected from the ocean, but develop over the slopes before expanding northwards towards the coast in the middle of the afternoon.

The third example (Figure 3) shows radar observations collected during the passage of tropical cyclone Dumazile over the Mascarene Archipelago in early March 2018. On 4 March (top panel), radar observations were collected within an outer rainband located about 700 km southeast of the TC center. Time series of vertical radar reflectivity profiles show numerous convective cells extending forward and backward up to an altitude of

15 km. On 6 March (bottom panel), radar data are collected within a cirriform region, at the back of the tropical cyclone. Vertical reflectivity profiles collected in this area (indicated by the red circle on the associated MODIS image) show a layer of cirrus clouds extending from 11 to 14 km that is characteristic of pre- and post-convective tropical cyclone regions. As shown by collocated high-resolution (6 min) rainfall measurements, brief but extremely strong rainfall occurred on 4 March around 09:00 UTC (up to 50 mmh^{-1}) and 10:15 UTC (up to 95 mmh^{-1}). These heavy precipitation periods are associated with data gaps in radar observations, which can also be observed in Figure 2a (between 8 and 11 UTC). These gaps reflect the aforementioned attenuation of the radar signal by strong precipitation, which principally occurs for instantaneous rain rates higher than 15 mm h^{-1} (Figures 2a and 3). According to the climatological study of [31], which provides a detailed analysis of the spatio-temporal variability of local rainfall patterns over Reunion Island, such situations are nevertheless uncommon in the Saint-Denis area, and not frequent enough to significantly impact the results that will be presented in the following.




Figure 2. Time series of radar reflectivity vertical profiles (dBZ) collected on (a) 26 March 2017 (summer) and (b) 1 August 2017 (winter). The black line in (a) shows collocated rainfall measurements every 6 min expressed in mmh^{-1} (scale to the left).

2.2. Other Datasets

Additional sources of data are also relied upon to describe the local and regional weather conditions prevailing over Reunion Island and the SWIO basin during the 2-year period of analysis:

- (i) In situ observations of relative humidity inferred from Saint-Denis' airport (20.88° S , 55.51° E) operational daily radiosounding data collected between 2014 and 2018;
- (ii) Time series of integrated water vapor (IWV) columns, inferred from the analysis of GNSS data collected from the IGN (Institut Geographique National) station STDE (20.88° S – 55.51° E) and the International GNSS System (IGS)' station REUN (21.2° S , 55.57° E). GNSS data are processed following the approach used by [34,35] to produce IWV data at various locations in the SWIO basin (a complete description of GNSS IWV retrieval and associated uncertainties can be found in the former papers).

- (iii) Reanalysis data extracted from European Center for Medium-Range Weather Forecast (ECMWF)'s 5th generation reanalysis (ERA5) are used to investigate the regional atmospheric conditions prevailing over the SWIO basin. ERA5 provides hourly estimates of numerous atmospheric, land and oceanic climate variables on a $0.25 \times 0.25^\circ$ horizontal grid [36]. Reanalysis data used in this study consist of wind, humidity, sea level pressure and vertical velocity fields extracted over the period 1990–2020.




Figure 3. Left panel: Satellite images (MODIS TERRA) of Tropical Cyclone Dumazile, recorded on 4 March 2018 (top, 9:00 UTC) and 6 March 2018 (bottom, 6:00 UTC). Right panel: corresponding time series of BASTA-REUNION vertical radar reflectivity profiles (dBZ, UTC) with superimposed collocated rainfall measurements every 6 min (expressed in mmh^{-1} , scale to the left). The red circle indicates the cirriform region sampled by the radar on 6 March.

3. Diurnal and Seasonal Variability of Cloudiness over Saint-Denis in 2017 and 2018

In the following, reflectivity data collected by BASTA-REUNION are used to investigate the vertical structure of winter and summer clouds over Saint-Denis. For simplicity, the four seasons investigated hereafter will be referred to as summer 2017 (November 2016–March 2017), winter 2017 (June–September 2017), summer 2018 (November 2017–March 2018) and winter 2018 (June–September 2018).

Diurnal Cycle

The seasonal composite analyses of radar reflectivity profiles aggregated over the two sampled summer and the two winter seasons are shown in Figure 4a,b, respectively.




Figure 4. Time series of cloud occurrence (% , normalized to 1) as a function of the time of the day (hours) and altitude (km) for (a) aggregated summer seasons 2017 and 2018, (b) aggregated winter seasons 2017 and 2018, (c) summer 2017, (d) winter 2017, (e) summer 2018, (f) winter 2018, (g) summer 2018 minus summer 2017 and (h) winter 2018 minus winter 2017. White lines near the altitudes 2, 5 and 10 km correspond to data gap layers of ~25–50 m resulting from the recombination of data collected in the different radar acquisition modes.

In summer (Figure 4a), aggregated radar observations show two cloud layers, associated with the diurnal cycle (leading to the formation of low-level cumulus clouds) and convective instability (leading to the development of thunderstorm systems in mid to late afternoon). In the lowest layer, which extends from ~0.5 to 3 km, the maximum cloud occurrence is observed between 08:00 UTC (12 LST) and 15:00 UTC (19 LST) at an altitude of 2 km, with a maximum probability of nearly 45% near 12 UTC (16 LST). The upper layer, comprised between ~6 and 15 km, corresponds to deep convective and ice clouds associated with local storms, as well as tropical storms and cyclones that passed nearby Reunion Island. The probability of occurrence is maximised between 10:00 UTC and 15:00 UTC (14–19 LST) and reaches up to 15% at an altitude of 10 km near 14 UTC (18 LST).

In winter (Figure 4b), one can still observe a layer of low-level clouds extending from approximately 0.5 to 2.5 km. The frequency of cloud occurrence is highest between 09:00 UTC (13 LST) and 14:00 UTC (18 LST), with a maximum probability of about 35% near 2km altitude at 11 UTC (15 LST). While the frequency of occurrence of low-level clouds appears similar during summer and winter periods, the depth of the low-level cloud layer is lower in winter (maximum height of 2.5 km vs. 3 km in summer). High-level clouds are also up to three times less frequent in winter (5% occurrence vs. 15% occurrence in summer) as a result of the stronger stability of the atmosphere during this season.

To further assess the diurnal variability of cloud cover from one year to the next, time series shown in Figure 4a,b were also individually aggregated for each of the four sampled seasons: summer 2017 (Figure 4c) and 2018 (Figure 4e), and winter 2017 (Figure 4d) and 2018 (Figure 4f). From a qualitative standpoint, one can observe strong similarities between both summers and both winters. However, a closer look at the field of differences (Figure 4g,h) shows important disparities from one year to the next. Hence, low-level cloudiness appears significantly higher and deeper in winter 2017 (Figure 4h) than during the following winter season, while significant differences could also be observed throughout the troposphere between the two summer seasons (Figure 4g), with significantly deeper cloud cover in 2018.

Mean vertical profiles of cloud occurrence inferred from the high-resolution BASTA-REUNION profiles collected during the four sampled seasons shown in Figure 4c–f are shown in Figure 5. During summer, the probability of occurrence shows much larger values in 2018 (solid blue line) at both low-level (frequency of cloud/precipitation occurrence below 3 km ranging from 20–22% vs. 15–20% in 2017) and mid-to-high level (frequency of cloud occurrence of 10% vs. 5% in 2017 at 6 km). These differences are even more striking in winter. In 2017 (red dashed line), low-level cloudiness is for instance twice as large as in 2018 (blue-dashed line, 20–22% vs. 10%) and actually bears strong resemblances with that observed during the previous summer (solid blue line). Furthermore, one can also observe large differences occurring at mid-levels. While the vertical cloud extension is more or less limited to 4 km in 2018, in good agreement with previous climatological studies of [30,31], clouds occur in much more significant proportions up to an altitude of 6 km and even beyond (frequency of occurrence of 1% at 8 km) in 2017. In the following, large-scale environmental conditions inferred from the analysis of ERA5 data are investigated in order to understand the possible origins of these differences.




Figure 5. Probability of cloud occurrence (% , normalized to 1) observed during summer (solid line) and winter (dashed line) 2017 (red) and 2018 (blue) as a function of height.

4. Analysis of Local and Large-Scale Environmental Conditions Prevailing in 2017 and 2018

4.1. IOSD Patterns

In the SWIO basin, the main climatic phenomenon that drives large-scale oceanic and atmospheric circulation is the Indian Ocean Subtropical Dipole (IOSD [37]). This climate

anomaly controls the structure of the ocean's surface layer, which, in turn, impacts the atmospheric circulation and rainfall at the regional scale. It is characterized by three distinct phases that are defined by a reference index computed from SST anomaly differences between the western and eastern parts of the basin: a neutral phase, corresponding to the mean state of the ocean, a negative phase and a positive phase. Positive events, which are characterized by above-normal SSTs in the southwestern part of the basin and below-normal SSTs off western Australia, generally start at the beginning of the Austral summer and die off in April/May.

The evolution of the IOSD index from 2013 to 2019 (Figure 6) shows that a particularly strong positive event occurred in early 2017. The latter started in November 2016, peaked in January (with a high index value) and finally died off in late March 2017 (index value of 0.2). However, one can observe that index values started to re-increase significantly at the beginning of the winter, with an average value of ~ 1 throughout the rest of the year, before finally collapsing in early 2018.




Figure 6. Averaged weekly (black) and quarterly (red) IOSD index between January 2013 and December 2019. Blue and green squares highlight index values during winter 2017 and 2018, respectively.

4.2. Large Scale Anomalies

Corresponding large-scale atmospheric and oceanic anomalies prevailing during summers (DJF) and winters (JJA) 2017 and 2018 are shown in Figures 7 and 8. In good agreement with the occurrence of a positive IOSD event, a positive SST anomaly, centered near 30° S and extending from the East African coast to $\sim 90^{\circ}$ E, can be observed over the SW part of the basin in summer 2017 (Figure 7a). Enhanced midlevel subsidence is also observed over Madagascar and the central part of the basin (Figure 7e), resulting in both positive pressure anomalies (Figure 7c) and dryer conditions in the mid-troposphere (as inferred from IWV anomalies, Figure 7g) in these areas. Over the Mascarenes and Reunion Island, these large-scale conditions result in a slightly positive SST anomaly, of nearly 0.5° C, and a below-normal (negative) humidity anomaly of $\sim 6\text{--}8\text{ kg m}^{-2}$.

During the next summer, a positive SST anomaly can still be observed in the southern part of the basin (Figure 7b). The latter is however shifted southeastwards with respect to summer 2017 and no significant surface pressure (Figure 7d) or mid-level vertical velocity (Figure 7f) anomalies are longer observed over the central part of the basin. A moist tongue extending northwestwards towards Madagascar could be observed, in good agreement with the location of the positive SST anomaly (Figure 7h). Over the Mascarene Archipelago, these large-scale conditions resulted in positive SST and moisture anomalies of up to 1° C and 8 kg m^{-2} , respectively.




Figure 7. (a,b) Sea Surface Temperature (SST), Mean Sea Level Pressure (MSLP), Vertical velocity at 500 hPa (W) and Integrated Water Vapour (IWV) anomalies in summer (DJF) 2017 (left panel) and 2018 (right panel), computed from ERA5 reanalysis data over the period 1990–2020.

During winter 2017 (Figure 8a), ERA5 data show SST anomalies of up to $1\text{ }^{\circ}\text{C}$ over the Mascarene Archipelago, likely inherited from the IOSD event that started earlier in the year. The Mascarene anticyclone also moved south-eastwards resulting in strong positive pressure anomalies over the central and southern parts of the basin as well as slightly below normal sea level pressures over the Mascarenes (Figure 8c). These anomalies are also accompanied by a decrease in the large-scale midlevel (500 hPa) subsidence over this area (Figure 8e), resulting in higher humidity content at 850 hPa (Figure 8g). While this positive SST anomaly is also present in winter 2018 (Figure 8b), the other atmospheric parameters (Figure 8d,f,h) are nevertheless globally in phase with the climatology.




Figure 8. (a,b) Sea Surface Temperature (SST), Mean Sea Level Pressure (MSLP), Vertical velocity at 500 hPa (W) and Integrated Water Vapour (IWV) anomalies in summer (DJF) 2017 (left panel) and 2018 (right panel), computed from ERA5 reanalysis data over the period 1990–2020. As in Figure 7, but for winter (JJA) 2017 and 2018.

4.3. Local Observations

Humidity observations inferred from the analysis of GNSS data collected at the IGS station of REUN (21.2° S, 55.57° E) and radio sounding data collected at Saint-Denis' airport (20.88° S, 55.51° E) are shown in Figure 9. According to time series of monthly averaged GNSS-derived IWV columns (Figure 9a), significantly dryer conditions occurred over Reunion Island during the first 4 months (Nov–March) of the wet season 2017, with differences of up to 12 kg m^{-2} ($+/-30\%$) compared to the same period one year after.

This tendency however inverts at the end of the summer, a period that corresponds to the dying of the 2017 positive IOSD event. GNSS-derived IWV observations also confirm that tropospheric humidity was significantly higher during winter 2017 (JJAS, blue line) than during winter 2018 (red line), with differences of up to 3.5 kg m^{-2} ($\pm 35\%$) between July and September. A strictly similar tendency (not shown) was also observed at other GNSS stations located along the west coast and central area of Reunion Island, indicating that these tropospheric humidity anomalies have impacted the whole island.




Figure 9. (a) Monthly mean of GNSS-derived IWV content at station REUN in 2017 (blue) and 2018 (red). (b) RS-derived relative humidity at Saint-Denis during summer (plain) and winter (dashed) 2017 (blue) and 2018 (red). The green line in (b) shows corresponding average RH profiles for the period 2014–2018.

According to seasonally averaged relative humidity data inferred from operational radio sounding observations in Saint-Denis (Figure 9b), absolute moisture conditions in summer 2018 (red solid line) were higher throughout the troposphere compared to summer 2017 (blue solid line), with an increase ranging from 10% at low levels to 100% at mid and high levels. The 5-y average (2014–2018) humidity profile (green solid line) also indicates that humidity conditions were significantly below (resp. above) normal in summer 2017 (resp. 2018), in good agreement with moisture anomaly patterns deduced from the analysis of ERA5 data (Figure 7g,h). In winter 2018, the seasonally averaged relative humidity profile (red dashed line) shows good agreement with the climatological mean (green dashed line) up to 6 km, and above normal relative humidity at higher levels. In 2017 (blue-dashed line), one can however observe a significant increase in the low-level relative humidity between 2 and 4 km altitude, ranging from +50% to +80% at a given level.

To further investigate relationships between cloudiness and humidity over Saint-Denis, Figure 10 shows hourly relative anomalies (difference between the hourly value and the daily average, divided by the daily average) of integrated water vapour (IWV, %), radar-derived low- (LLC, 1–3 km) and mid-to-high-level (HLC, 5–10 km) cloudiness (%) as well as insolation, during summer (left panel) and winter (right panel) 2018 (as GNSS

observations collocated with BASTA-REUNION only started to be available in January 2018, we can only compare cloud occurrence and IWV for this year).




Figure 10. Diurnal cycle of high level (HLC) and low-level (LLC) clouds, Integrated water vapour (IWV) and insolation at Saint-Denis in 2018. Top panel: low-(1–3 km, plain) and mid-(5–10 km, dashed) level cloud anomaly (%) during (a) summer 2018 and (b) winter 2018. Bottom panel: Diurnal cycles of GNSS-derived (STDE station) IWV anomaly (%), black and insolation (hours, orange) during (c) summer 2018 and (d) winter 2018.

The ground starts to heat up in the early morning (3 UTC/7 LST) with sunrise (Figure 10c,d), which increases the air temperature and initiates thermal instability in the boundary layer. As water vapour generated by evapotranspiration starts to condense, a positive cloudiness anomaly appears near 7:30 UTC (11:30 LST) at both low- and mid-to-high levels. A positive IWV anomaly, which remains positive throughout the daytime, can also be observed at nearly the same hour. In both summer (Figure 10c) and winter (Figure 10d) this positive moisture anomaly is maximum near 12 UTC (16 LST) and reaches about 4.5% ($\sim 2 \text{ kg m}^{-2}$ in summer and 1.06 kg m^{-2} in winter). The time of this maximum likely reflects the existence of a sea breeze regime, which is strongest in mid-afternoon, when the difference in temperature between the land and ocean is highest. This IWV anomaly maximum also corresponds to that of the observed maximum in low-level cloudiness anomaly, which reaches up to 60% in summer (Figure 10a) and 50% in winter. This agreement suggests that the sea breeze regime may act to reinforce low-level cloud formation along the slopes through the advection of additional moisture originating from the ocean towards the island.

5. Discussion and Conclusions

The permanent deployment, in 2016, of a 95 GHz vertically pointing cloud radar BASTA in Reunion Island is an important achievement in order to reinforce cloud observations and atmospheric science research capabilities in the SWIO.

Data collected by this radar during its two first years of operation show a strong seasonal cycle with significant variation of cloudiness between summers and winters. During Austral summer, a bimodal mode characterized by two cloud layers extending respectively from 1–3 km (maximum daily frequency of cloud occurrence of 45%) and 8–12 km (maximum daily frequency of cloud occurrence of 15%) is observed. During Austral winter, cloudiness is mostly limited to the low levels (maximum daily probability of occurrence of about 35%) due to both the northward migration of the ITCZ and the

strong trade wind inversion prevailing at this time of the year (which further limits vertical cloud extension). These results are globally in line with those obtained by [19] over the tropical island of Barbados (using ground-based cloud radar data collected over a similar period of two years), which have also shown the existence of a bimodal mode (in the vertical) in summer that disappears in winter due to the circulation shifts accompanying the seasonal migration of the ITCZ. As in [19], we also observed a slight variability in the lower troposphere cloudiness across the seasons on a diurnal scale. In summer, medium and high clouds associated with convective developments over the island's relief show a maximum occurrence between 10 and 15UTC (i.e., between 14 and 19 LST), while low-level clouds are maximised between 8 and 15 UTC (12 and 19 LST) in summer and 9 and 14 UTC (13 and 18 LST) in winter. As in Barbados, this variability appears somewhat more important in summer (maximum value of cloud occurrence of 45% at a height of 2 km at both locations) than during winter (25% in Barbados vs. 35% in Reunion Island at a height of 2 km). The diurnal cycle of the cloudiness is also strongly correlated with the diurnal cycle of IWV regardless of the season and the cloud layer considered.

The comparative analysis of seasonal cloudiness observations collected in 2017 and 2018 also shows strong differences from one year to the next, with significantly higher (resp. lower) cloud cover in winter (resp. summer) 2017 than in 2018. The analysis of ERA5 reanalysis data and local humidity measurements derived from radio soundings and GNSS observations suggests that these differences could be attributed to a different large-scale context between the two years. In summer 2017 a strong positive SST anomaly, which formed in relationship with a positive phase of the subtropical Indian Ocean dipole (SIOD), could be observed south of Madagascar and was accompanied by positive (resp. negative) anomalies of subsidence (resp. humidity) over the center part of the basin. This large-scale subsidence, and associated dry air, acted to limit the vertical extension of clouds and likely explains the significant differences in cloud cover observed between the two summers. The two winter seasons also show important differences in terms of the large-scale environment. In 2017, a zonal dipole of large-scale subsidence anomaly was observed, while conditions were closer to normal in 2018. The eastward shift of high subsidence values resulting from the presence of this dipole resulted in a wet anomaly over the western part of the basin in winter 2017 that was further amplified by a positive SST anomaly over Reunion Island (increased evaporation). As a consequence, the vertical structure of the cloudiness and the probability of cloud occurrence in the lower half of the troposphere were surprisingly similar throughout the year 2017.

According to regional and local humidity data, one can hypothesize that the 2017 IOSD event significantly impacted moisture conditions over Reunion Island throughout the year 2017, resulting in lower (resp. higher) than normal humidity conditions at low- and mid-level in summer (resp. winter). These distinct moisture conditions may have accounted for the large differences observed in terms of cloudiness between Austral summers 2017 and 2018 (Figure 4). The thermodynamic structure of the atmosphere over Reunion Island was also more unstable during Austral winter 2017 than during Austral winter 2018, thus favouring low-level cloud development, which eventually happens to overshoot the trade wind inversion. In this regard, an exceptional and extremely intense storm, which locally produced more than 105 mm of rainfall in one hour in the northeast part of the island, occurred on 29 and 30 August 2017. BASTA-REUNION reflectivity measurements collected at Saint-Denis during this event (Figure 11) show convective cells with vertical development above 7 km altitude, which is extremely uncommon in this season. Instantaneous rainfall data collected next to the radar on 30 August around 1:30 LST confirms that particularly intense rainfall (up to 70 mmh^{-1}) occurred over Saint-Denis even though the rainfall maxima was observed about 15 km SE of the radar. Additionally, note the extinction of the radar signal at the same hour, as already observed in Figures 2 and 3.




Figure 11. As in Figure 2, but for data collected between 29 August (10 UTC/14 LST) and 30 August (5 UTC/9 LST) 2017.

Decreased moisture at low levels during summer 2017 can also account for the limited development of deep local convective events with respect to 2018. The observed differences in terms of cloudiness were also exacerbated by the significant cyclonic activity that occurred over the Mascarene Archipelago during the wet season 2018. Between January and April 2018 (see Figure 1 in [38]), four tropical cyclones grazed the island (TC Berguitta and Fakir) or passed in its immediate vicinity (TC Ava and Dumazile)—which drained a large amount of mid-to-high level clouds over Reunion Island—whereas only two tropical storms (TS Carlos and Fernando), which passed more than 200 km away from the island, occurred in 2017.

Cloud radar observations are key to investigate the properties and variability of tropical clouds, to evaluate cloud fraction forecasted by NWP systems [39] and derived from space-borne measurements, as well as to evaluate climate model performance [40]. Although limited in time, the preliminary analysis of the 2-y dataset of cloud radar observations analysed in this study already allows us to better understand the various and complex processes that drive cloudiness in the tropical island of the SWIO such as Reunion Island. More efforts are nevertheless obviously needed in order to thoroughly characterize the properties of the clouds in this tropical area as well as to fully exploit the rare observations provided by research cloud radars in the Southern hemisphere.

In this regard, observations of clouds and precipitation in Reunion Island will continue to be reinforced in the frame of ACTRIS through the deployment, in late 2021, of a second, volumetric (i.e., 360° scanning), BASTA cloud radar at the Maïdo observatory. A transportable polarimetric X-band radar, aiming at complementing observations provided by Reunion Island’s two operational S-band weather radar systems, will also be installed at Saint-Joseph (southern part of the island) in September 2021, in the frame of the Interreg-V Indian Ocean research project ESPOIRS (“Etude des Systèmes Précipitant de l’Océan Indien par Radars et Satellites” [41]). The deployment of this additional cloud radar, together with the enhancement of the ground-based weather radar network, will allow for further, long-term, studies of the seasonal and inter-annual variability of tropical clouds, but also to benefit from multi-frequency radar observations to better understand the formation and life cycle of clouds and precipitation over this unique tropical area. This new setup will thus permit us to study all sorts of cloud patterns, ranging from orographic, cirrus and low-to-mid level trade clouds to deep convective storms associated with tropical cyclones passing nearby. Altogether, these instruments will also contribute to make OPAR and Reunion Island a precious satellite validation testbed in the southern hemisphere for Earth observation programs, such as the upcoming EARTHCARE [42] space mission.

Author Contributions: Conceptualization J.D. (Jonathan Durand) and O.B.; data acquisition and processing J.D. (Julien Delanoë), J.D. (Jonathan Durand) and O.B.; writing—original draft preparation, J.D. (Jonathan Durand), E.L., F.B. and O.B.; writing—review and editing, all co-authors; supervision, O.B.; project administration, O.B.; funding acquisition, O.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the European Union, the Regional Council of Reunion Island and the French state under FEDER projects “BASTA-Reunion” and “ESIEC-RUN” and INTERREG-V Indian Ocean 2014–2020 project “ReNovRisk-Cyclone and Precipitation”.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data used in this study are available on request from the corresponding author.

Acknowledgments: Authors are grateful to the engineers of the Laboratoire Atmosphère, Milieux et Observations Spatiales (LATMOS) for their help in the setup and operation of the BASTA-REUNION radar.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Liou, K.-N. Influence of Cirrus Clouds on Weather and Climate Processes: A Global Perspective. *Mon. Weather. Rev.* **1986**, *114*, 1167–1199. [CrossRef]
- Hansen, J.; Sato, M.; Ruedy, R. Radiative forcing and climate response. *J. Geophys. Res. Space Phys.* **1997**, *102*, 6831–6864. [CrossRef]
- Klein, S.A.; Hall, A.; Norris, J.R.; Pincus, R. Low-Cloud Feedbacks from Cloud-Controlling Factors: A Review. *Surv. Geophys.* **2017**, *38*, 1307–1329. [CrossRef]
- Bony, S. Marine boundary layer clouds at the heart of tropical cloud feedback uncertainties in climate models. *Geophys. Res. Lett.* **2005**, *32*, 20806. [CrossRef]
- Stephens, G.L. Cloud Feedbacks in the Climate System: A Critical Review. *J. Clim.* **2005**, *18*, 237–273. [CrossRef]
- Nuijens, L.; Stevens, B.; Siebesma, A.P. The Environment of Precipitating Shallow Cumulus Convection. *J. Atmos. Sci.* **2009**, *66*, 1962–1979. [CrossRef]
- Madden, R.; Julian, P. Detection of a 40-50-day oscillation of the zonal wind in the tropical Pacific. *J. Atmos. Sci.* **1971**, *28*, 702–708. [CrossRef]
- Madden, R.; Julian, P. Observations of the 40–50-day tropical oscillation. A review. *Mon. Wea. Rev.* **1994**, *122*, 814–837. [CrossRef]
- Battaglia, A.; Kollias, P.; Dhillon, R.; Lamer, K.; Khairoutdinov, M.; Watters, D. Mind the gap—Part 2: Improving quantitative estimates of cloud and rain water path in oceanic warm rain using spaceborne radars. *Atmos. Meas. Tech.* **2020**, *13*, 4865–4883. [CrossRef]
- Marchand, R.; Mace, G.G.; Ackerman, T.; Stephens, G. Hydrometeor Detection Using Cloudsat—An Earth-Orbiting 94-GHz Cloud Radar. *J. Atmos. Ocean. Technol.* **2008**, *25*, 519–533. [CrossRef]
- Sassen, K.; Wang, Z. Classifying clouds around the globe with the CloudSat radar: 1-year of results. *Geophys. Res. Lett.* **2008**, *35*, 04805. [CrossRef]
- Lamer, K.; Kollias, P.; Battaglia, A.; Preval, S. Mind the gap—Part 1: Accurately locating warm marine boundary layer clouds and precipitation using spaceborne radars. *Atmos. Meas. Tech.* **2020**, *13*, 2363–2379. [CrossRef]
- Kollias, P.; Clothiaux, E.E.; Miller, M.A.; Albrecht, B.A.; Stephens, G.L.; Ackerman, T.P. Millimeter-Wavelength Radars: New Frontier in Atmospheric Cloud and Precipitation Research. *Bull. Am. Meteorol. Soc.* **2007**, *88*, 1608–1624. [CrossRef]
- Long, C.N.; Mather, J.H.; Ackerman, T.P. The ARM Tropical Western Pacific (TWP) Sites. *Meteorol. Monogr.* **2016**, *57*, 7. [CrossRef]
- Thorsen, T.J.; Fu, Q.; Comstock, J. Cloud effects on radiative heating rate profiles over Darwin using ARM and A-train radar/lidar observations. *J. Geophys. Res. Atmos.* **2013**, *118*, 5637–5654. [CrossRef]
- Liu, Z.; Marchand, R.; Ackerman, T. A comparison of observations in the tropical western Pacific from ground-based and satellite millimeter-wavelength cloud radars. *J. Geophys. Res. Space Phys.* **2010**, *115*. [CrossRef]
- Bouniol, D.; Couvreur, F.; Kamsu-Tamo, P.; Leplay, M.; Guichard, F.; Favot, F.; O’Connor, E.J. Diurnal and Seasonal Cycles of Cloud Occurrences, Types, and Radiative Impact over West Africa. *J. Appl. Meteorol. Clim.* **2012**, *51*, 534–553. [CrossRef]
- Sukanya, P.; Kalapureddy, M.C.R. Cloud radar observations of multi-scale variability of cloud vertical structure associated with Indian summer monsoon over a tropical location. *Clim. Dyn.* **2021**, *56*, 1055–1081. [CrossRef]
- Stevens, B.; Farrell, D.; Hirsch, L.; Jansen, F.; Nuijens, L.; Serikov, I.; Brüggemann, B.; Forde, M.; Linné, H.; Lonitz, K.; et al. The Barbados Cloud Observatory: Anchoring Investigations of Clouds and Circulation on the Edge of the ITCZ. *Bull. Am. Meteorol. Soc.* **2016**, *97*, 787–801. [CrossRef]
- Stevens, B.D.; Bony, S.; Farrell, D.; Ament, F. EUREC4A, submitted to Earth. *Sys. Sci. Data.* 2021. Available online: <https://essd.copernicus.org/preprints/essd-2021-18/> (accessed on 29 May 2021).

21. Brueck, M.; Nuijens, L.; Stevens, B. On the Seasonal and Synoptic Time-Scale Variability of the North Atlantic Trade Wind Region and Its Low-Level Clouds. *J. Atmos. Sci.* **2015**, *72*, 1428–1446. [[CrossRef](#)]
22. Kollias, P.; Bharadwaj, N.; Clothiaux, E.E.; Lamer, K.; Oue, M.; Hardin, J.; Isom, B.; Lindenmaier, I.; Matthews, A.; Luke, E.; et al. The ARM Radar Network: At the Leading Edge of Cloud and Precipitation Observations. *Bull. Am. Meteorol. Soc.* **2020**, *101*, E588–E607. [[CrossRef](#)]
23. Baray, J.-L.; Courcoux, Y.; Keckhut, P.; Portafaix, T.; Tulet, P.; Cammas, J.-P.; Hauchecorne, A.; Godin Beekmann, S.; De Mazière, M.; Hermans, C.; et al. Maïdo observatory: A new high-altitude station facility at Reunion Island (21° S, 55° E) for long-term atmospheric remote sensing and in situ measurements. *Atmos. Meas. Tech.* **2013**, *6*, 2865–2877. [[CrossRef](#)]
24. Delanoë, J.; Protat, A.; Vinson, J.-P.; Brett, W.; Caudoux, C.; Bertrand, F.; Du Chatelet, J.P.; Hallali, R.; Barthes, L.; Haeffelin, M.; et al. BASTA: A 95-GHz FMCW Doppler Radar for Cloud and Fog Studies. *J. Atmos. Ocean. Technol.* **2016**, *33*, 1023–1038. [[CrossRef](#)]
25. Bergemann, M.; Jakob, C. How important is tropospheric humidity for coastal rainfall in the tropics? *Geophys. Res. Lett.* **2016**, *43*, 5860–5868. [[CrossRef](#)]
26. Dupont, J.-C.; Haeffelin, M.; Wærsted, E.; Delanoë, J.; Renard, J.-B.; Preissler, J.; O’Dowd, C. Evaluation of Fog and Low Stratus Cloud Microphysical Properties Derived from In Situ Sensor, Cloud Radar and SYRSOC Algorithm. *Atmosphere* **2018**, *9*, 169. [[CrossRef](#)]
27. Quetelard, H.; Bessemoulin, P.; Cerveny, R.S.; Peterson, T.C.; Burton, A.; Boodhoo, Y. Extreme Weather: World-Record Rainfalls During Tropical Cyclone Gamede. *Bull. Am. Meteorol. Soc.* **2009**, *90*, 603–608. [[CrossRef](#)]
28. Taupin, F.G.; Bessafi, M.; Baldy, S.; Bremaud, P.J. Tropospheric ozone above the southwestern Indian Ocean is strongly linked to dynamical conditions prevailing in the tropics. *J. Geophys. Res. Space Phys.* **1999**, *104*, 8057–8066. [[CrossRef](#)]
29. Brémaud, P.; Taupin, F. Cloud influence on ozone diurnal cycle in the marine boundary layer at Réunion Island. *Atmos. Res.* **1998**, *47–48*, 285–298. [[CrossRef](#)]
30. Lesouëf, D.; Gheusi, F.; Delmas, R.; Escobar, J. Numerical simulations of local circulations and pollution transport over Reunion Island. *Ann. Geophys.* **2011**, *29*, 53–69. [[CrossRef](#)]
31. Réchou, A.; Flores, O.; Jumaux, G.; Duflot, V.; Bousquet, O.; Pouppeville, C.; Bonnardot, F. Spatio-temporal variability of rainfall in a high tropical island: Patterns and large-scale drivers in Réunion Island. *Q. J. R. Meteorol. Soc.* **2019**, *145*, 893–909. [[CrossRef](#)]
32. Rechou, A.; Rao, T.N.; Bousquet, O.; Plu, M.; Decoupes, R. Properties of rainfall in a tropical volcanic island deduced from UHF wind profiler measurements. *Atmos. Meas. Tech.* **2014**, *7*, 409–418. [[CrossRef](#)]
33. Jumeaux, G.; Quetelard, H.; Roy, D. Atlas Climatique de La Réunion de Météo-France, Météo-France technical report 2011. (in French). Available online: <http://www.meteofrance.fr/publications/nos-collections/climat-outr-mer/atlas-climatique-de-la-reunion> (accessed on 29 May 2021).
34. Lees, E.; Bousquet, O.; Roy, D.; Leclair, J. Analysis of diurnal to seasonal variability of integrated water vapour in the South Indian Ocean basin using ground-based GNSS and 5th generation ECMWF reanalysis (ERA5) data. *Q. J. R. Meteorol.* **2021**, *147*, 229–248. [[CrossRef](#)]
35. Bousquet, O.; Lees, E.; Durand, J.; Peltier, A.; Duret, A.; Mekies, D.; Boissier, P.; Donal, T.; Fleischer-Dogley, F.; Zakariasy, L. Densification of the Ground-Based GNSS Observation Network in the Southwest Indian Ocean: Current Status, Perspectives, and Examples of Applications in Meteorology and Geodesy. *Front. Earth Sci.* **2020**, *8*. [[CrossRef](#)]
36. Hersbach, H.; Bell, B.; Berrisford, P.; Hirahara, S.; Horanyi, A.; Muñoz-Sabater, J.; Nicolas, J.; Peubey, C.; Radu, R.; Schepers, D.; et al. The ERA5 global reanalysis. *Q. J. R. Meteorol. Soc.* **2020**, *146*, 1999–2049. [[CrossRef](#)]
37. Behera, S.; Yamagata, T. Subtropical SST dipole events in the southern Indian Ocean. *Geophys. Res. Lett.* **2001**, *28*, 327–330. [[CrossRef](#)]
38. Bousquet, O.; Barbary, D.; Bielli, S.; Kebir, S.; Raynaud, L.; Malardel, S.; Faure, G. An evaluation of tropical cyclone forecast in the Southwest Indian Ocean basin with AROME-Indian Ocean convection-permitting numerical weather predicting system. *Atmos. Sci. Lett.* **2019**, *21*, 950. [[CrossRef](#)]
39. Hogan, R.J.; O’Connor, E.J.; Illingworth, A.J. Verification of cloud-fraction forecasts. *Q. J. R. Meteorol. Soc.* **2009**, *135*, 1494–1511. [[CrossRef](#)]
40. Protat, A.; Bouniol, D.; Delanoë, J.; May, P.T.; Plana-Fattori, A.; Hasson, A.; Görsdorf, U.; Heymsfield, A.J. Assessment of Cloudsat Reflectivity Measurements and Ice Cloud Properties Using Ground-Based and Airborne Cloud Radar Observations. *J. Atmos. Ocean. Technol.* **2009**, *26*, 1717–1741. [[CrossRef](#)]
41. Bousquet, O.; Barruol, G.; Cordier, E.; Barthe, C.; Bielli, S.; Calmer, R.; Rindraharisaona, E.; Roberts, G.; Tulet, P.; Amelie, V.; et al. Impact of Tropical Cyclones on Inhabited Areas of the SWIO Basin at Present and Future Horizons. Part 1: Overview and Observing Component of the Research Project RENOVRIK-CYCLONE. *Atmosphere* **2021**, *12*, 544. [[CrossRef](#)]
42. Illingworth, A.J.; Barker, H.W.; Beljaars, A.; Ceccaldi, M.; Chepfer, H.; Clerbaux, N.; Cole, J.; Delanoë, J.; Domènech, C.; Donovan, D.P.; et al. The EarthCARE Satellite: The Next Step Forward in Global Measurements of Clouds, Aerosols, Precipitation, and Radiation. *Bull. Am. Meteorol. Soc.* **2015**, *96*, 1311–1332. [[CrossRef](#)]

36. Cesca, S.; Letort, J.; Razafindrakoto, H.N.T.; Heimann, S.; Rivalta, E.; Isken, M.P.; Nikkhoo, M.; Passarelli, L.; Petersen, G.M.; Cotton, F.; et al. Drainage of a deep magma reservoir near Mayotte inferred from seismicity and deformation. *Nat. Geosci.* **2020**, *13*, 87–93. [[CrossRef](#)]
37. Davy, C.; Barruol, G.; Fontaine, F.R.; Sigloch, K.; Stutzmann, E. Tracking major storms from microseismic and hydroacoustic observations on the seafloor. *Geophys. Res. Lett.* **2014**, *41*, 8825–8831. [[CrossRef](#)]
38. Davy, C.; Barruol, G.; Fontaine, F.; Cordier, E. Analyses of extreme swell events on La Réunion Island from microseismic noise. *Geophys. J. Int.* **2016**, *207*, 1767–1782. [[CrossRef](#)]
39. Barruol, G.; Davy, C.; Fontaine, F.R.; Schlindwein, V.; Sigloch, K. Monitoring austral and cyclonic swells in the “Iles Eparses” (Mozambique channel) from microseismic noise. *Acta Oecologica* **2016**, *72*, 120–128. [[CrossRef](#)]
40. Rindraharisaona, E.J.; Cordier, E.; Barruol, G.; Fontaine, F.R.; Singh, M. Assessing swells in La Réunion Island from terrestrial seismic observations, oceanographic records and offshore wave models. *Geophys. J. Int.* **2020**, *221*, 1883–1895. [[CrossRef](#)]
41. Rindraharisaona, E.J.; Barruol, G.; Cordier, E.; Fontaine, F. and Gonzales, A.. Inferring cyclone signatures in the South-West of Indian Ocean from microseismic noise recorded on Réunion island. *Atmosphere* **2021**, *12*, 488. [[CrossRef](#)]
42. Bousquet, O.; Barbary, D.; Bielli, S.; Kebir, S.; Raynaud, L.; Malardel, S.; Faure, G. An evaluation of tropical cyclone forecast in the Southwest Indian Ocean basin with AROME-Indian Ocean convection-permitting numerical weather predicting system. *Atmos. Sci. Lett.* **2020**, *21*, e950. [[CrossRef](#)]
43. Hsiao, L.-F.; Chen, D.-S.; Hong, J.-S.; Yeh, T.-C.; Fong, C.-T. Improvement of the Numerical Tropical Cyclone Prediction System at the Central Weather Bureau of Taiwan: TWRF (Typhoon WRF). *Atmosphere* **2020**, *11*, 657. [[CrossRef](#)]
44. Takeuchi, Y. An Introduction of Advanced Technology for Tropical Cyclone Observation, Analysis and Forecast in JMA. *Trop. Cyclone Res. Rev.* **2019**, *7*, 153–163. [[CrossRef](#)]
45. Mehra, A.; Tallapragada, V.; Zhang, Z.; Liu, B.; Zhu, L.; Wang, W.; Kim, H.S. Advancing the State of the Art in Operational Tropical Cyclone Fore-casting at NCEP. *Trop. Cyclone Res. Rev.* **2018**, *7*, 51–56. [[CrossRef](#)]
46. Davidson, N.E.; Xiao, Y.; Ma, Y.; Weber, H.C.; Sun, X.; Rikus, L.J.; Kepert, J.D.; Steinle, P.X.; Dietachmayer, G.S.; Lok, C.C.F.; et al. ACCESS-TC: Vortex Specification, 4DVAR Initialization, Verification, and Structure Diagnostics. *Mon. Weather Rev.* **2014**, *142*, 1265–1289. [[CrossRef](#)]
47. Bender, M.A.; Ginis, I. Real-case simulations of hurricane-ocean interaction using a high-resolution coupled model: Effects on hurricane intensity. *Mon. Weather Rev.* **2000**, *128*, 917–946. [[CrossRef](#)]
48. Black, W.J.; Dickey, T.D. Observations and analyses of upper ocean responses to tropical storms and hurricanes in the vicinity of Bermuda. *J. Geophys. Res. Space Phys.* **2008**, *113*, 08009. [[CrossRef](#)]
49. Price, J.F. Upper ocean response to a hurricane. *J. Phys. Ocean.* **1981**, *11*, 153–175. [[CrossRef](#)]
50. Jullien, S.; Marchesiello, P.; Menkes, C.E.; Lefèvre, J.; Jourdain, N.C.; Samson, G.; Lengaigne, M. Ocean feedback to tropical cyclones: Climatology and processes. *Clim. Dyn.* **2014**, *43*, 2831–2854. [[CrossRef](#)]
51. Smith, R.K.; Montgomery, M.T.; Van Sang, N. Tropical cyclone spin-up revisited. *Q. J. R. Meteorol. Soc.* **2009**, *135*, 1321–1335. [[CrossRef](#)]
52. Andreas, E.L. Spray Stress Revisited. *J. Phys. Oceanogr.* **2004**, *34*, 1429–1440. [[CrossRef](#)]
53. Doyle, J.D. Coupled Atmosphere–Ocean Wave Simulations under High Wind Conditions. *Mon. Weather Rev.* **2002**, *130*, 3087–3099. [[CrossRef](#)]
54. Andreas, E.L.; Edson, J.B.; Monahan, E.C.; Rouault, M.P.; Smith, S.D. The spray contribution to net evaporation from the sea: A review of recent progress. *Boundary-Layer Meteorol.* **1995**, *72*, 3–52. [[CrossRef](#)]
55. Wang, Y.; Kepert, J.D.; Holland, G.J. The Effect of Sea Spray Evaporation on Tropical Cyclone Boundary Layer Structure and Intensity*. *Mon. Weather Rev.* **2001**, *129*, 2481–2500. [[CrossRef](#)]
56. Liu, B.; Liu, H.; Xie, L.; Guan, C.; Zhao, D. A Coupled Atmosphere–Wave–Ocean Modeling System: Simulation of the Intensity of an Idealized Tropical Cyclone. *Mon. Weather Rev.* **2011**, *139*, 132–152. [[CrossRef](#)]
57. Richter, D.H.; Stern, D.P. Evidence of spray-mediated air-sea enthalpy flux within tropical cyclones. *Geophys. Res. Lett.* **2014**, *41*, 2997–3003. [[CrossRef](#)]
58. Andreas, E.L.; Mahrt, L.; Vickers, D. An improved bulk air-sea surface flux algorithm, including spray-mediated transfer. *Q. J. R. Meteorol. Soc.* **2015**, *141*, 642–654. [[CrossRef](#)]
59. Ovadnevaite, J.; Manders, A.; De Leeuw, G.; Ceburnis, D.; Monahan, C.; Partanen, A.-I.; Korhonen, H.; O’Dowd, C.D. A sea spray aerosol flux parameterization encapsulating wave state. *Atmos. Chem. Phys. Discuss.* **2014**, *14*, 1837–1852. [[CrossRef](#)]
60. Willoughby, H.; Jin, H.-L.; Lord, S.J.; Piotrowicz, J.M. Hurricane structure and evolution as simulated by an ax-isymmetric, nonhydrostatic numerical model. *J. Atmos. Sci.* **1984**, *41*, 1169–1186. [[CrossRef](#)]
61. Lord, S.J.; Willoughby, H.E.; Piotrowicz, J.M. Role of a parameterized ice-phase microphysics in an axisymmet-ric, nonhydrostatic tropical cyclone model. *J. Atmos. Sci.* **1984**, *41*, 2836–2848. [[CrossRef](#)]
62. Wang, Y. An Explicit Simulation of Tropical Cyclones with a Triply Nested Movable Mesh Primitive Equation Model: TCM3. Part II: Model Refinements and Sensitivity to Cloud Microphysics Parameterization*. *Mon. Weather Rev.* **2002**, *130*, 3022–3036. [[CrossRef](#)]
63. Zhu, T.; Zhang, D.-L. Numerical Simulation of Hurricane Bonnie (1998). Part II: Sensitivity to Varying Cloud Microphysical Processes. *J. Atmos. Sci.* **2006**, *63*, 109–126. [[CrossRef](#)]

64. Li, J.; Wang, G.; Lin, W.; He, Q.; Feng, Y.; Mao, J. Cloud-scale simulation study of Typhoon Hagupit (2008) Part II: Impact of cloud microphysical latent heat processes on typhoon intensity. *Atmos. Res.* **2013**, *120*, 202–215. [[CrossRef](#)]
65. Hoarau, T.; Barthe, C.; Tulet, P.; Claeys, M.; Pinty, J.-P.; Bousquet, O.; Delanoë, J.; Vié, B. Impact of the Generation and Activation of Sea Salt Aerosols on the Evolution of Tropical Cyclone Dumile. *J. Geophys. Res. Atmos.* **2018**, *123*, 8813–8831. [[CrossRef](#)]
66. Bielli, S.; Barthe, C.; Bousquet, O.; Tulet, P.; Pianezze, J. The effect of atmosphere-ocean coupling on the structure and intensity of tropical cyclone Bejisa observed in the southwest Indian ocean. *Atmosphere* **2021**. submitted to this Special Issue.
67. Thompson, C.; Barthe, C.; Bielli, S.; Tulet, P.; Pianezze, J. Projected Characteristic Changes of a Typical Tropical Cyclone under Climate Change in the South West Indian Ocean. *Atmosphere* **2021**, *12*, 232, (this Special Issue). [[CrossRef](#)]
68. Ferrario, F.; Beck, M.W.; Storlazzi, C.D.; Micheli, F.; Shepard, C.C.; Airolidi, L. The effectiveness of coral reefs for coastal hazard risk reduction and adaptation. *Nat. Commun.* **2014**, *5*, 3794. [[CrossRef](#)]
69. Lowe, R.J.; Falter, J.L.; Bandet, M.D.; Pawlak, G.; Atkinson, M.J.; Monismith, S.G.; Koseff, J.R. Spectral wave dissipation over a barrier reef. *J. Geophys. Res. Space Phys.* **2005**, *110*, 04001. [[CrossRef](#)]
70. van Dongeren, A.; Storlazzi, C.D.; Quataert, E.; Pearson, S. Wave dynamics and flooding on low-lying tropical reef-lined coasts. In Proceedings of the Coastal Dynamics, Helsingor, Denmark, 12–16 June 2017; pp. 654–664.
71. Pomeroy, A.; Lowe, R.J.; Symonds, G.; Van Dongeren, A.; Moore, C. The dynamics of infragravity wave transformation over a fringing reef. *J. Geophys. Res. Space Phys.* **2012**, *117*, 11022. [[CrossRef](#)]
72. Baldock, T.; Golshani, A.; Callaghan, D.; Saunders, M.; Mumby, P. Impact of sea-level rise and coral mortality on the wave dynamics and wave forces on barrier reefs. *Mar. Pollut. Bull.* **2014**, *83*, 155–164. [[CrossRef](#)] [[PubMed](#)]
73. Nortek. Wave Measurements Using the PUV Method, TN-19. Doc. No. N4000-720, Nortek AS. 2002. Available online: <http://www.nortek-es.com/lib/technical-notes/puv-wave-measurement> (accessed on 20 March 2021).
74. Hom-Ma, M.; Horikawa, K.; Komori, S. Response Characteristics of Underwater Wave Gauge. *Coast. Eng. Jpn.* **1966**, *9*, 45–54. [[CrossRef](#)]
75. Pomeroy, A.W.M.; Lowe, R.J.; Ghisalberti, M.; Winter, G.; Storlazzi, C.; Cuttler, M. Spatial Variability of Sediment Transport Processes Over Intratidal and Subtidal Timescales Within a Fringing Coral Reef System. *J. Geophys. Res. Earth Surf.* **2018**, *123*, 1013–1034. [[CrossRef](#)]
76. Smithers, S.G.; Hoeke, R.K. Geomorphological impacts of high-latitude storm waves on low-latitude reef is-lands—observations of the December 2008 event on Nukutoa, Takuu, Papua New Guinea. *Geomorphology* **2014**, *222*, 106–121. [[CrossRef](#)]
77. Hoeke, R.K.; McInnes, K.L.; O’Grady, J.G. Wind and Wave Setup Contributions to Extreme Sea Levels at a Tropical High Island: A Stochastic Cyclone Simulation Study for Apia, Samoa. *J. Mar. Sci. Eng.* **2015**, *3*, 1117–1135. [[CrossRef](#)]
78. Friedrich, A.; Krüger, F.; Klinge, K. Ocean-generated microseismic noise located with the Gräfenberg array. *J. Seismol.* **1998**, *2*, 47–64. [[CrossRef](#)]
79. Longuet-Higgins, M.S. A theory of the origin of the microseisms. *Phil. Trans. Roy. Soc.* **1950**, *243*, 1–35.
80. Arduin, F.; Gualtieri, L.; Stutzmann, E. How ocean waves rock the Earth: Two mechanisms explain microseisms with periods 3 to 300 s. *Geophys. Res. Lett.* **2015**, *42*, 765–772. [[CrossRef](#)]
81. Barruol, G.; Reymond, D.; Fontaine, F.R.; Hyvernaud, O.; Maurer, V.; Maamaatuaiahutapu, K. Characterizing swells in the southern Pacific from seismic and infrasonic noise analyses. *Geophys. J. Int.* **2006**, *164*, 516–542. [[CrossRef](#)]
82. Cessaro, R.K. Sources of primary and secondary microseisms. *Bull. Seismol. Soc. Am.* **1994**, *84*, 142–148.
83. Hasselmann, K. A statistical analysis of the generation of microseisms. *Rev. Geophys.* **1963**, *1*, 177–210. [[CrossRef](#)]
84. Arduin, F.; Stutzmann, E.; Schimmel, M.; Mangeney, A. Ocean wave sources of seismic noise. *J. Geophys. Res. Space Phys.* **2011**, *116*, 09004. [[CrossRef](#)]
85. Essen, H.-H.; Krüger, F.; Dahm, T.; Grevemeyer, I. On the generation of secondary microseisms observed in northern and central Europe. *J. Geophys. Res. Space Phys.* **2003**, *108*, 2506. [[CrossRef](#)]
86. Obrebski, M.; Arduin, F.; Stutzmann, E.; Schimmel, M. How moderate sea states can generate loud seismic noise in the deep ocean. *Geophys. Res. Lett.* **2012**, *39*, 11601. [[CrossRef](#)]
87. Davy, C.; Stutzmann, E.; Barruol, G.; Fontaine, F.; Schimmel, M. Sources of secondary microseisms in the Indian Ocean. *Geophys. J. Int.* **2015**, *202*, 1180–1189. [[CrossRef](#)]
88. Reading, A.M.; Koper, K.D.; Gal, M.; Graham, L.S.; Tkalčić, H.; Hemer, M.A. Dominant seismic noise sources in the Southern Ocean and West Pacific, 2000–2012, recorded at the Warramunga Seismic Array, Australia. *Geophys. Res. Lett.* **2014**, *41*, 3455–3463. [[CrossRef](#)]
89. Beucler, É.; Mocquet, A.; Schimmel, M.; Chevrot, S.; Quillard, O.; Vergne, J.; Sylvander, M. Observation of deep water microseisms in the North Atlantic Ocean using tide modulations. *Geophys. Res. Lett.* **2015**, *42*, 316–322. [[CrossRef](#)]
90. Bromirski, P.D. The near-coastal microseism spectrum: Spatial and temporal wave climate relationships. *J. Geophys. Res. Space Phys.* **2002**, *107*, ESE 5-1–ESE 5-20. [[CrossRef](#)]
91. Bromirski, P.D.; Duennebier, F.K.; Stephen, R.A. Mid-ocean microseisms. *Geochem. Geophys. Geosystems* **2005**, *6*, Q04009. [[CrossRef](#)]
92. Chevrot, S.; Sylvander, M.; Benahmed, S.; Ponsolles, C.; Lefèvre, J.M.; Paradis, D. Source locations of secondary microseisms in western Europe: Evidence for both coastal and pelagic sources. *J. Geophys. Res. Space Phys.* **2007**, *112*, B11301. [[CrossRef](#)]
93. Koper, K.D.; Buriaciu, R. The fine structure of double-frequency microseisms recorded by seismometers in North America. *J. Geophys. Res.* **2015**, *120*, 1677–1691. [[CrossRef](#)]

94. Tolman, H.L.; Chalikov, D.V. Source terms in a third-generation wind wave model. *J. Phys. Oceanogr.* **1996**, *26*, 2497–2518. [[CrossRef](#)]
95. Tolman, H.L. Distributed-memory concepts in the wave model WAVEWATCH III. *Parallel Comput.* **2002**, *28*, 35–52. [[CrossRef](#)]
96. Perez, J.; Menendez, M.; Losada, I.J. GOW2: A global wave hindcast for coastal applications. *Coast. Eng.* **2017**, *124*, 1–11. [[CrossRef](#)]
97. Hasselmann, K.; Barnett, T.; Bouws, E.; Carlson, H.; Cartwright, D.E.; Eake, K. *Measurements of Wind-Wave Growth and Swell Decay during the Joint North Sea Wave Project (JONSWAP)*; Deutsches Hydrographisches Institute: Hamburg, Germany, 1973.
98. Harris, L.R.; Nel, R.; Oosthuizen, H.; Meyer, M.; Kotze, D.; Anders, D.; McCue, S.; Bachoo, S. Managing conflicts between economic activities and threatened migratory marine species toward creating a multiobjective blue economy. *Conserv. Biol.* **2017**, *32*, 411–423. [[CrossRef](#)]
99. Swart, N.C.; Lutjeharms, J.R.E.; Ridderinkhof, H.; De Ruijter, W.P.M. Observed characteristics of Mozambique Channel eddies. *Geophys. Res. Lett.* **2010**, *115*, C09006. [[CrossRef](#)]
100. Lutjeharms. *The Agulhas Current*; Springer: Berlin/Heidelberg, Germany, 2006. [[CrossRef](#)]
101. Lellouche, J.-M.; Greiner, E.; Le Galloudec, O.; Garric, G.; Regnier, C.; Drevillon, M.; Benkiran, M.; Testut, C.-E.; Bourdalle-Badie, R.; Gasparin, F.; et al. Recent updates to the Copernicus Marine Service global ocean monitoring and forecasting real-time 1/12° high-resolution system. *Ocean Sci.* **2018**, *14*, 1093–1126. [[CrossRef](#)]
102. Cuyppers, Y.; Le Vaillant, X.; Bourruet-Aubertot, P.; Vialard, J.; McPhaden, M.J. Tropical storm-induced near-inertial internal waves during the Cirene experiment: Energy fluxes and impact on vertical mixing. *J. Geophys. Res. Oceans* **2013**, *118*, 358–380. [[CrossRef](#)]
103. Vialard, J.; Duvel, J.; McPhaden, M.; Bourruet-Aubertot, P.; Ward, P.; Key, E.; Bourras, D.; Weller, R.; Minnett, P.J.; Weill, A.; et al. Cirene: Air Sea Interactions in the Sey-chelles-Chagos thermocline ridge region. *Bull. Am. Met. Soc.* **2009**, *90*, 45–61. [[CrossRef](#)]
104. Jaimes, B.; Shay, L.K. Mixed Layer Cooling in Mesoscale Oceanic Eddies during Hurricanes Katrina and Rita. *Mon. Weather Rev.* **2009**, *137*, 4188–4207. [[CrossRef](#)]
105. Montégut, C.D.B.; Madec, G.; Fischer, A.S.; Lazar, A.; Iudicone, D. Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology. *J. Geophys. Res. Space Phys.* **2004**, *109*, 1–20. [[CrossRef](#)]
106. Holte, J.; Talley, L. A New Algorithm for Finding Mixed Layer Depths with Applications to Argo Data and Subantarctic Mode Water Formation*. *J. Atmos. Ocean. Technol.* **2009**, *26*, 1920–1939. [[CrossRef](#)]
107. Tomczak, M.; Godfrey, J.S. *Regional Oceanography—An Introduction*; Butler & Tanner: London, UK, 2001.
108. Pickard, G.L.; Emery, W.J. *Descriptive Physical Oceanography*, 5th ed.; Butterworth-Heinemann: Oxford, UK, 1990.
109. Roberts, G.; Burnet, F.; Barrau, S.; Medina, P.; Durand, P.; Gavard, M. A drone over the oceans: The Miriad project. *Météorologie* **2017**, *98*, 9–10. (In French) [[CrossRef](#)]
110. Bevis, M.; Businger, S.; Herring, T.A.; Rocken, C.; Anthes, R.A.; Ware, R.H. GPS meteorology: Remote sensing of atmospheric water vapor using the global positioning system. *J. Geophys. Res. Space Phys.* **1992**, *97*, 15787–15801. [[CrossRef](#)]
111. Beutler, G.; Rothacher, M.; Schaer, S.; Springer, T.; Kouba, J.; Neilan, R. The International GPS Service (IGS): An interdisciplinary service in support of Earth sciences. *Adv. Space Res.* **1999**, *23*, 631–653. [[CrossRef](#)]
112. Guerova, G.; Jones, J.; Douša, J.; Dick, G.; de Haan, S.; Pottiaux, E.; Bock, O.; Pacione, R.; Elgered, G.; Vedel, H.; et al. Review of the state of the art and future prospects of the ground-based GNSS meteorology in Europe. *Atmos. Meas. Tech.* **2016**, *9*, 5385–5406. [[CrossRef](#)]
113. Dvorak, V.F. Tropical Cyclone Intensity Analysis and Forecasting from Satellite Imagery. *Mon. Weather Rev.* **1975**, *103*, 420–430. [[CrossRef](#)]
114. Brennan, M.J.; Hennon, C.C.; Knabb, R.D. The Operational Use of QuikSCAT Ocean Surface Vector Winds at the National Hurricane Center. *Weather Forecast* **2009**, *24*, 621–645. [[CrossRef](#)]
115. Chou, K.-H.; Wu, C.-C.; Lin, S.-Z. Assessment of the ASCAT wind error characteristics by global dropwindsonde observations. *J. Geophys. Res. Atmos.* **2013**, *118*, 9011–9021. [[CrossRef](#)]
116. Mai, M.; Zhang, B.; Li, X.; Hwang, P.A.; Zhang, J.A. Application of AMSR-E and AMSR2 Low-Frequency Channel Brightness Temperature Data for Hurricane Wind Retrievals. *IEEE Trans. Geosci. Remote. Sens.* **2016**, *54*, 4501–4512. [[CrossRef](#)]
117. Zhang, L.; Yin, X.-B.; Shi, H.-Q.; Wang, Z.-Z. Hurricane Wind Speed Estimation Using WindSat 6 and 10 GHz Brightness Temperatures. *Remote. Sens.* **2016**, *8*, 721. [[CrossRef](#)]
118. Reul, N.; Chapron, B.; Zabolotskikh, E.; Donlon, C.; Mouche, A.; Tenerelli, J.; Collard, F.; Piolle, J.-F.; Fore, A.; Yueh, S.; et al. A New Generation of Tropical Cyclone Size Measurements from Space. *Bull. Am. Meteorol. Soc.* **2017**, *98*, 2367–2385. [[CrossRef](#)]
119. Monserrat, O.; Crosetto, M.; Luzi, G. A review of ground-based SAR interferometry for deformation measurement. *ISPRS J. Photogramm. Remote. Sens.* **2014**, *93*, 40–48. [[CrossRef](#)]
120. Fu, L.; Holt, B. *SEASAT Views Oceans and Sea Ice with Synthetic Aperture Radar*; NASA JPL Publ.: La Cañada Flintridge, CA, USA, 1982; pp. 81–120. Available online: https://earth.esa.int/documents/10174/1020083/Seasat_views_Oceans_Sea_Ice_with_SAR.pdf (accessed on 20 March 2021).
121. Li, X.; Zhang, J.A.; Yang, X.; Pichel, W.G.; DeMaria, M.; Long, D.; Li, Z. Tropical Cyclone Morphology from Spaceborne Synthetic Aperture Radar. *Bull. Am. Meteorol. Soc.* **2013**, *94*, 215–230. [[CrossRef](#)]
122. Young, G.; Sikora, T.; Winstead, N. Mesoscale Near-Surface Wind Speed Variability Mapping with Synthetic Aperture Radar. *Sensors* **2008**, *8*, 7012–7034. [[CrossRef](#)]

123. Zhang, B.; Perrie, W. Cross-Polarized Synthetic Aperture Radar: A New Potential Measurement Technique for Hurricanes. *Bull. Am. Meteorol. Soc.* **2012**, *93*, 531–541. [[CrossRef](#)]
124. Mouche, A.; Chapron, B.; Knaff, J.; Zhao, Y.; Zhang, B.; Combot, C. Copolarized and Cross-Polarized SAR Measurements for High-Resolution Description of Major Hurricane Wind Structures: Application to Irma Category 5 Hurricane. *J. Geophys. Res. Oceans* **2019**, *124*, 3905–3922. [[CrossRef](#)]
125. Duong, Q.-P.; Langlade, S.; Payan, C.; Husson, R.; Mouche, A.; Malardel, S. C-band SAR Winds for Tropical Cyclone monitoring and forecast in the South-West Indian Ocean. *Atmosphere* **2021**. accepted pending minor revisions (this Special Issue).
126. Mouche, A.; Chapron, B.; Zhang, B.; Husson, R. Combined co- and cross-polarized SAR measurements under extreme wind conditions. *IEEE Xplore IEEE Trans. Geosci. Remote Sens.* **2017**, *55*, 6476–6755.
127. Sitkowski, M.; Kossin, J.P.; Rozoff, C.M. Intensity and Structure Changes during Hurricane Eyewall Replacement Cycles. *Mon. Weather Rev.* **2011**, *139*, 3829–3847. [[CrossRef](#)]
128. Willoughby, H.; Clos, J.; Shoreibah, M.L. Concentric Eye Walls, Secondary Wind Maxima, and The Evolution of the Hurricane vortex. *J. Atmos. Sci.* **1982**, *39*, 395–411. [[CrossRef](#)]
129. Madec, G.; Bourdallé-Badie, R.; Chanuta, J. *Nemo Ocean Engine. Notes From the Pôle de Modélisation*; Institut Pierre-Simon Laplace: Guyancourt, France, 2019.
130. Barbary, D.; Leroux, M.-D.; Bousquet, O. The orographic effect of Reunion Island on tropical cyclone track and intensity. *Atmos. Sci. Lett.* **2019**, *20*, e882. [[CrossRef](#)]

